VERSION 1
MAR 19, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bthnnj5e

Protocol Citation: Karla LH Feijs 2021. Calcium phosphate transfection mammalian cells.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bthnnj5e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
This our routine protocol for calcium phosphate transfections.

Created: Mar 19, 2021

Last Modified: Mar 19, 2021

PROTOCOL integer ID:
48398

🌐 Calcium phosphate transfection mammalian cells V.1

Karla LH
Feijs¹

¹Institute of Biochemistry and Molecular Biology, RWTH Aachen University Hospital

FeZa lab

Karla LH Feijs
Institute of Biochemistry and Molecular Biology, RWTH Aachen...

DISCLAIMER

This protocol is a mix of the protocols I've been taught during my master internship in René Bernards lab and during my PhD in Bernhard Lüscher's lab.

ABSTRACT

This is our calcium phosphate transfection protocol. It always works with HEK293T cells, for other lines the transfection efficiency depends on their source: some HeLa and U2OS clones can be transfected using CaPO₄, however most cannot. We mostly use this method to transfect HEK293T cells for lentivirus production.

MATERIALS

10x HBS-buffer:

40 gr. NaCl

25 gr. Hepes

1.85 gr. KCl

0.63 gr. Na₂HPO₄·H₂O

500ml ddH₂O

Store in aliquots of 50mL at -20°C.

1x HBS-buffer:

Dilute 50mL 10x HBS-buffer in a total volume of 450mL ddH₂O and adjust pH to **6.95** with NaOH. Adjust the volume to 500mL with ddH₂O.

Sterilize by passage through a 22mm-filter. Store in 50 mL aliquots at 4°C.

2.5M CaCl₂:

18.4 gr. CaCl₂·2H₂O in 50 mL ddH₂O.

Sterilize by passage through a 22mm-filter. Store at 4°C.

10x HEPES

5 gr. KCl

83 gr. NaCl

24 gr. HEPES

- or - 26 gr. Na-HEPES

Add ±800 mL ddH₂O.

Adjust the pH to **7.3**, adjust volume to 1L and autoclave.

1x HEPES

Dilute 50mL in 450mL sterile ddH₂O to make 1x HEPES. Or dilute in 450 mL ddH₂O and autoclave.

BEFORE START INSTRUCTIONS

The CRUCIAL part in this protocol is the adjustment of the pH of the HEBS buffer EXACTLY. We do this at room temperature with room temperature water.

1 Day 1

Seed cells: density depends on cell line and dish used. For example in a 10cm dish seed 0.8x10⁶ HeLa or 1.5 x10⁶ HEK293T cells; in a 6-well plate seed 0.3 x10⁶ HEK293T cells or 0.15x10⁶ HeLa cells.

2 Day 2

Observe the cells under the microscope.

Prepare the transfection mix. The amounts given here are for a 10cm dish and can be adjusted to other dishes and plates.

Prepare in an eppendorf tube:

20µg plasmid DNA

Add 950µl HEBS buffer

Add 50µl CaCl₂

Flick the tube gently to thoroughly mix and incubate at room temperature for approximately 20 minutes.

We rarely use a full 20µg of DNA encoding for something: some times as little as 2 µg of the total is encoding for a protein of interest, whereas the rest is an empty vector. This amount of DNA is required to efficiently form precipitates.

- 3 Add the mixture drop-wise onto the cells, gently agitate the dish to evenly distribute the mixture and place the cells back in the incubator.
- 4 After 5-6 hours, remove the media, add warm HEPES buffer and incubate 5-10 minutes. Remove the HEPES and add fresh full DMEM. The precipitates are visible under a microscope and should disappear during the incubation with HEPES. Replace the cells in the incubator.

5 Day 3-4

Process your cells 24 to 48 hours after transfection, incubation time depending on experimental needs and set-up. If it does not interfere with the experiment, 5% of the transfected plasmid DNA can be a GFP-encoding plasmid to monitor transfection efficiency.