NOV 20, 2020

🌐 Calculating Colony-forming Units

Jiri Hulcr¹, You Li¹

¹University of Florida

Protocols Bark Beetle Mycobiome

Bark Beetle Mycobiome Research Coordination Network

ABSTRACT

This protocol describes how to estimate the number of colony-forming units.

This protocol is part of the Bark Beetle Mycobiome (BBM) Research Coordination Network. For more information on the BBM international network: Hulcr J, Barnes I, De Beer ZW, Duong TA, Gazis R, Johnson AJ, Jusino MA, Kasson MT, Li Y, Lynch S, Mayers C, Musvuugwa T, Roets F, Seltmann KC, Six D, Vanderpool D, & Villari C. 2020. Bark beetle mycobiome: collaboratively defined research priorities on a widespread insect-fungus symbiosis. *Symbiosis* 81: 101–113 <https://doi.org/10.1007/s13199-020-00686-9>.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bnuvmew6

Document Citation: Jiri Hulcr, You Li 2020. Calculating Colony-forming Units. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bnuvmew6>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Oct 23, 2020

Last Modified: Nov 20, 2020

DOCUMENT integer ID:
43637

To estimate the initial number of colony-forming units in a mycangium, use the technique of serial dilution when plating. After colonies grow, multiply the number of colonies on the plate by the inverse of the initial dilution factor.

1. Prepare two tubes per each sample, label them “0.1” and “0.01”, fill each with 500 ul of water or PBS.
2. Suspend mycangium in the tube “0.1” and vortex.
3. Plate **50 ul** of the suspension on media. Record the plate as “0.1 dilution” in [PLATES].[note] in the Isolations database.
4. Transfer **50 ul** of the initial suspension to the second tube (“0.01”) and vortex.

5. Plate **50 ul** of the second suspension on **second** media plate, and record that plate as "0.01 dilution".
6. Plate **5 ul** of the second suspension on a **third** plate, and record it as "0.001 dilution".