

Dec 13, 2018

Version 1

Cambridge Fetal Thymus Dissociation V.1

DOI

dx.doi.org/10.17504/protocols.io.wedfba6

Jongeun Park¹

¹Wellcome Sanger Institute

Human Cell Atlas Metho...



Jongeun Park

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.wedfba6>

Protocol Citation: Jongeun Park 2018. Cambridge Fetal Thymus Dissociation. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.wedfba6>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

We use this protocol and it's working

Created: December 13, 2018

Last Modified: December 13, 2018

Protocol Integer ID: 18597

Keywords: Fetal, Thymus, Spleen, protocol for fetal thymus dissociation, fetal thymus dissociation, cambridge fetal thymus dissociation, other fetal tissue, lymphoid organ, thymus, digestion, larger tissue size, multiple rounds of digestion, tissue, account of larger tissue size

Abstract

This is protocol for fetal thymus dissociation. (Can be applied to other fetal tissues)

Can be also adapted to adult thymus or lymphoid organs. (Volume of solution needs to be adjusted to take account of larger tissue size, also consider multiple rounds of digestion in this case)

Attachments



Cambridge Fetal thym...

15KB

Troubleshooting

