MAY 08, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.ydtfs6n

External link:
<https://mmpc.org/shared/document.aspx?id=271&docType=Protocol>

Protocol Citation: Henri Brunengraber 2019. Case - Intraperitoneal Glucose Tolerance Test. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.ydtfs6n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 20, 2019

Last Modified: May 08, 2019

PROTOCOL integer ID:
 20627

Case - Intraperitoneal Glucose Tolerance Test

Henri Brunengraber¹

¹Case Western Reserve University

Mouse Metabolic Phenotyping Centers
 Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

This is the standard protocol for most routine glucose tolerance testing. It is performed on awake mice, fasted for 18 hr (overnight) or 6 hours.

MATERIALS

MATERIALS

- ⊗ D-Glucose stock 25 mg/ml in saline Sigma Aldrich
- ⊗ 1cc syringe 26g BD Biosciences
- ⊗ Clotisol Amazon Catalog #T212002
- ⊗ AlphaTrak2 Contributed by users
- ⊗ Glucometer Alphasat Alphasat Catalog #cfs185-m0285

Reagents and Materials:

Reagent/Material	Quantity Required	Vendor	Stock Number
D-Glucose stock, 25 mg/ml in saline	2gm/kg body weight	Sigma	
1cc syringe 26g	1/Mouse	BD	
Glucometer Alphasat	1		cfs185-m0285
AlphaTrak2 Glucose strips	5 per mouse		
Clotisol	100ul	Amazon	T212002

Note:

Sigma-Aldrich, [RRID:SCR_008988](#)
 BD Biosciences, [RRID:SCR_013311](#)

Keywords: Intraperitoneal
Glucose Tolerance Test

- 1 Fast mice overnight or for 6 hours. Remove mouse from cage and put into a clean cage with water and no food (5:00PM). Next day begin GTT by 9:00AM.
- 2 Weigh each mouse and record weight.
- 3 Insert glucose strip into glucometer and check that the code matches for the strip being used.
- 4 Take fasting blood glucose by snipping the tail and putting a drop on the glucose strip (already in glucometer). Record fasting glucose as time 0'. Dip tail into clotisol to clot blood.
- 5 Calculate the amount of glucose needed for a concentration of 2gm glucose/kg body weight. Record the volume for each mouse.
- 6 Inject each mouse 3' apart for time 0. Repeat #4 at 15min, 30 min, 60 min and 120 min after IP injection of glucose.
- 7 Do not take extra blood for insulin measurements. This causes too much stress. Use a separate group of mice for collecting blood after glucose injection for insulin measurements.