SEP 08, 2017

cell cultures

PLOS One

yuejun wang¹

¹Department of Prosthodontics, Peking University School and Hospital of Stomatology

Yuejun Wang

ABSTRACT

Teeth were first disinfected with 75% ethanol and then washed with phosphate-buffered saline. Briefly, PDLSCs were separated from periodontal ligament in the middle one-third of the root. Subsequently, MSCs were digested in a solution of 3 mg/mL collagenase type I (Worthington Biochemical Corp., Lakewood, NJ, USA) and 4 mg/mL dispase (Roche Diagnostics Corp., Indianapolis, IN, USA) for 1 h at 37°C. Single-cell suspensions were obtained by cell passage through a 70-µm strainer (Falcon, BD Labware, Franklin Lakes, NJ, USA). MSCs were grown in a humidified, 5% CO₂ incubator at 37°C in DMEM alpha modified Eagle's medium (Invitrogen, Carlsbad, CA, USA), supplemented with 15% fetal bovine serum (FBS; Invitrogen, Carlsbad, CA, USA), 2 mmol/L glutamine, 100 U/mL penicillin and 100 µg/mL streptomycin (Invitrogen, Carlsbad, CA, USA). The culture medium was changed every 3 days. MSCs at passages 3-5 were used in subsequent experiments. Human embryonic kidney 293T cells were maintained in complete DMEM with 10% FBS, 100 U/mL penicillin and 100 µg/mL streptomycin.

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.iegcbbw](https://doi.org/10.17504/protocols.io.iegcbbw)

External link:

<https://doi.org/10.1371/journal.pone.0184182>

Protocol Citation: yuejun wang 2017. cell cultures.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.iegcbbw>

MANUSCRIPT CITATION:

Wang Y, Liu Y, Fan Z, Liu D, Wang F, Zhou Y (2017) IGFBP2 enhances adipogenic differentiation potentials of mesenchymal stem cells from Wharton's jelly of the umbilical cord *via* JNK and Akt signaling pathways. PLoS ONE 12(8): e0184182. doi:

[10.1371/journal.pone.0184182](https://doi.org/10.1371/journal.pone.0184182)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jun 12, 2017

Last Modified: Feb 02, 2018

PROTOCOL integer ID:
6312

Teeth preparation

- 1 Teeth were first disinfected with 75% ethanol and then washed with phosphate-buffered saline.

Tissue preparation

- 2 PDLSCs were separated from periodontal ligament in the middle one-third of the root.

cell digestion

- 3 MSCs were digested in a solution of 3 mg/mL collagenase type I (Worthington Biochemical Corp., Lakewood, NJ, USA) and 4 mg/mL dispase (Roche Diagnostics Corp., Indianapolis, IN, USA) for 1 h at 37°C.

Single-cell suspension

- 4 Single-cell suspensions were obtained by cell passage through a 70-µm strainer (Falcon, BD Labware, Franklin Lakes, NJ, USA).

cell culture

- 5 MSCs were grown in a humidified, 5% CO₂ incubator at 37°C in DMEM alpha modified Eagle's medium (Invitrogen, Carlsbad, CA, USA), supplemented with 15% fetal bovine serum (FBS; Invitrogen, Carlsbad, CA, USA), 2 mmol/L glutamine, 100 U/mL penicillin and 100 µg/mL streptomycin (Invitrogen, Carlsbad, CA, USA). The culture medium was changed every 3 days.

MSCs at passages 3-5 were used in subsequent experiments.