APR 15, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.zzwf77e

Protocol Citation: George Lykotrafitis 2019. Centrifuge the whole blood to separate red blood cells. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.zzwf77e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol in our group and it is working.

Created: Apr 11, 2019

Last Modified: Apr 15, 2019

PROTOCOL integer ID:
22294

Centrifuge the whole blood to separate red blood cells

George Lykotrafitis¹

¹University of Connecticut

George Lykotrafitis

ABSTRACT

Centrifuge the whole blood in order to obtain the red blood cells.

MATERIALS

Human whole blood, centrifuge, centrifuge tubes.

- 1 Centrifuge the whole blood using the parameters: 500 ×g, 4 °C, and 5 min.

- 2 Remove the plasma and buffy coat.
- 3 Add the same volume of Alsever's solution as the remaining blood components.
- 4 Centrifuge again using the parameters: 500 ×g, 4 °C, and 5 min.
- 5 Remove the top layer.
- 6 Add the same volume of Alsever's solution as the remaining blood components.
- 7 Centrifuge a third time using the parameters: 500 ×g, 4 °C, and 5 min.
- 8 Remove the top layer.
- 9 Add the same volume of Alsever's solution as the remaining blood component - red blood cells, and store at 4 °C for up to 7 days.