OCT 16, 2019

🌐 Chitin binding assay using chitin magnetic beads (NEB)

Robert Hooftman¹

¹Wageningen University

iGEM Wageningen 2019

Robert Hooftman
Wageningen University

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8anhdsde

Protocol Citation: Robert Hooftman 2019. Chitin binding assay using chitin magnetic beads (NEB). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.8anhdsde>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Oct 16, 2019

Last Modified: Oct 16, 2019

PROTOCOL integer ID:
28718

MATERIALS

MATERIALS

⊗ Chitin Magnetic Beads - 5 ml **New England Biolabs Catalog #E8036S**

STEP MATERIALS

⊗ Chitin Magnetic Beads - 5 ml **New England Biolabs Catalog #E8036S**

1 Make the following buffer:

1 X CBD Column Binding Buffer

Tris-HCl	20 mM Adjust pH to 8 using HCl	(2,42 g/L)
NaCl	500 mM	(29,22 g/L)
EDTA	1 mM	(0,292 g/L)
Tween-20	0.05% v/v	500 ul/ 1 L

Gently mix the beads solution and resuspend thoroughly.

Chitin Magnetic Beads - 5 ml **New England Biolabs Catalog #E8036S**

- 2 Aliquot 50 ul of beads solution into a 2 ml microfuge tube
- 3 Apply a magnet for 30 seconds to pull the beads to one side. Remove the excess liquid 3s
- 4 Wash the beads 3 times with 500 ul of 1x CBD Column Binding Buffer. Apply magnet and remove the liquid.
- 5 Load 200-500 ul of cell supernatant with the beads.
Note: when making dilutions of the supernatant, dilute in CBD Column Binding Buffer.
- 6 Mix thoroughly and incubate for 1 hour at 4°C at constant agitation. 1h
- 7 Apply magnet for 30 seconds and remove supernatant with a pipette. Save the supernatant. 3s

- 8** Wash beads 3 times with 500 ul of 1x CBD Column Binding Buffer. Apply magnet for 30 seconds and pipette off liquid. Save the wash fractions. 3s
- 9** To remove proteins from the beads, boil beads for 5 minutes in 50 ul SDS-Page sample buffer at 95 °C 5m
- 10** Run fractions on a protein gel