MAR 07, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.ywvfxe6

Protocol Citation: Nicolas M Schmelling 2019. Chlorophyll Extraction in Cyanobacteria. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.ywvfxe6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Mar 07, 2019

Last Modified: Mar 07, 2019

PROTOCOL integer ID:
 21173

Chlorophyll Extraction in Cyanobacteria

Forked from [Chlorophyll Extraction in Cyanobacteria](#)

Nicolas M Schmelling¹

¹Institute of Synthetic Microbiology Heinrich Heine University

M4455 - Synthetische Biologie und Biotechnologie

Sebastian ST Triesch

Institute for Synthetic Microbiology, HHU Düsseldorf

ABSTRACT

This protocol should be used for chlorophyll extraction in cyanobacteria. The equation for calculating the exact chlorophyll content can be found at the end of this document.

You might want to measure the optical density (OD) of your cyanobacteria culture at 750 nm. Use BG11 medium or water as the reference solution. You need the OD of your culture to normalize the chlorophyll concentration to the number of cyanobacteria.

Calculate chlorophyll content (adapted from [Lichtenthaler 1978](#))

$$\text{Chl } [\mu\text{g/ml}] = \text{OD}_{665\text{nm}} \times 13.9 [\mu\text{g/ml}] \times \text{dilution factor of culture}$$

You can take less than 1 ml, but note the dilution factor for the calculation later on, e.g. :

1 ml sample = dilution factor of 1
 500 µl sample = dilution factor of 2
 100 µl sample = dilution factor of 10

- 1 Take **1 ml sample** of your cyanobacteria culture and spin it down at **14,000 rpm** for **5 min**.

⌚ 00:05:00

- 2 **Discard 0.9 ml** of the **supernatant**. **Resuspend** the **pellet** in the **remaining 100 µl**.
- 3 **Add 0.9 ml** of **100% methanol** to the sample and **mix thoroughly** by vortexing.
- 4 Incubate the samples in the **dark** for **30 min** at **4 °C** in the fridge.
00:30:00
- 5 Spin down samples again at **14,000 rpm** for **5 min**.
00:05:00
- 6 **Transfer supernatant** into a cuvette and measure the **extinction** at **665 nm**. Use **90% methanol** as the **reference** solution.