MAY 18, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bfd9ji96

Protocol Citation: New England Biolabs 2020. Cleavage of the Fusion Protein (TEV Protease). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bfd9ji96>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Apr 21, 2020

Last Modified: May 18, 2020

PROTOCOL integer ID:
 36001

Keywords: TEV, protease, cleavage

Cleavage of the Fusion Protein (TEV Protease)

New England Biolabs¹

¹New England Biolabs

New England Biolabs (NEB)

Tech. support phone: +1(800)632-7799 email: info@neb.com

New England Biolabs

New England Biolabs

ABSTRACT

TEV Protease, also known as Tobacco Etch Virus (TEV) Protease, is a highly specific cysteine protease that recognizes the amino-acid sequence Glu-Asn-Leu-Tyr-Phe-Gln-(Gly/Ser) and cleaves between the Gln and Gly/Ser residues. It is often used for the removal of affinity purification tags such as maltose-binding protein (MBP) or poly-histidine from fusion proteins. TEV Protease has a 7xHis-tag for easy removal from a reaction using nickel affinity resins and has been engineered to improve thermal stability and decrease autolysis.

GUIDELINES

Affinity tags, such as Maltose Binding Protein (MBP) or polyhistidine (His), are essential tools for the production of recombinant proteins. MBP is known to significantly enhance the solubility of many proteins, resulting in higher yield. Whereas, His-tagging is widely employed for the purification of recombinant target proteins via immobilized metal affinity chromatography (IMAC). The His-tag advantages include small tag size and high affinity and specificity of poly-His tag binding to divalent metals at neutral pH. Despite these important advantages, it is often preferred to remove the affinity tag following purification in order to isolate the target protein. Although chemical and enzymatic methods can be used, proteolytic enzymes, such as TEV Protease (NEB #[P8112](#)) and Factor Xa (NEB #[P8010](#)) are the preferred method for cleavage of fusion proteins at designed cleavage sites.

MATERIALS

MATERIALS

TEV protease **New England Biolabs Catalog #P8112S**

SAFETY WARNINGS

Please refer to the Safety Data Sheets (SDS) for health and environmental hazards.

BEFORE START INSTRUCTIONS

MBP and the target protein are fused by a polylinker containing a TEV protease recognition site for easy removal of the MBP-tag. One unit of TEV Protease will cleave approximately 2 μg of fusion protein. Cleavage should be carried out in 1X TEV Protease Reaction Buffer or in Amylose column elution buffer supplemented with DTT to a final concentration of 1 mM. Depending on the particular fusion protein, the amount of TEV Protease can be adjusted to get an acceptable rate of cleavage.

1 If necessary, concentrate the fusion protein to **at least** $[M]$ 0.5 milligram per milliliter (mg/mL) .

2 Perform a pilot experiment with a small portion of your protein. For example:

2.1 Combine flask 15 μg fusion protein and H₂O to make a **total reaction volume of 45 μL** .

2.2 Add flask 5 μL TEV Protease Reaction Buffer (10X) to make a **50 μL total reaction volume**.

2.3 Add flask 1 μL TEV Protease .

2.4 In a separate tube, combine flask 5 μg fusion protein ,
 flask 5 μL TEV Protease Reaction Buffer (10X) and H₂O to a volume of **50 μL** . **Do not add TEV Protease (control sample).**

2.5 Incubate reaction and control sample for 01:00:00 , 03:00:00 , and 08:00:00 at 30 °C (an additional reaction can be made and incubated for 24 hours at 4°C).

2.6 Take 10 µL of reaction(s) at the indicated times above and add 5 µL SDS-PAGE Sample Buffer (3X) . Take 10 µL control sample and add 5 µL SDS-PAGE Sample Buffer (3X) after 8 hours (or longest incubation time).

2.7 Incubate the SDS-PAGE samples for 00:03:00 - 00:05:00 at 70 °C - 100 °C .

2.8 Analyze them by SDS-PAGE.

3 Scale up the pilot experiment linearly for the amount of the fusion protein to be cleaved. Save at least a small sample of the uncut fusion as a negative control.

4 Check for complete cleavage by SDS-PAGE.

5 TEV Protease and the cleaved MBP contain polyhistidine tags at their N-termini. They can be removed from the cleavage reaction by immobilized metal affinity chromatography, such as Nickel or Cobalt resin, thereby isolating the target protein in the flow through.