OCT 18, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.utcewiw

Protocol Citation: Ralitsa R Madsen 2018. Coating of plates with Geltrex, for human iPSC culture.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.utcewiw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 18, 2018

Last Modified: Oct 18, 2018

PROTOCOL integer ID:
 16964

🌐 Coating of plates with Geltrex, for human iPSC culture

Ralitsa R Madsen¹

¹University of Edinburgh

Ralitsa R Madsen

University College London, University of Edinburgh, Universi...

GUIDELINES

Geltrex Catalog NO: A1413302 (5 ml from ThermoFischerScientific). Thin Gel Method (non-gelling) for hESC Applications. From:
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3157883/>

SAFETY WARNINGS

When working with volumes of Geltrex <5ml, aliquot to appropriate required working volumes and store at -20 to -80°C. Avoid multiple freeze/thaw cycles.

BEFORE START INSTRUCTIONS

Thaw Geltrex™ Basement Membrane Matrix at 2-8°C overnight. Refrigerator temperatures may vary; therefore, thaw extract on ice in a refrigerator. Basement Membrane Matrix gels in 5-10 minutes above 15°C; therefore when working from a full 5 ml vial, it is unnecessary to keep it on ice if used within 5 minutes and the environmental temperature does not exceed 25°C. It is also unnecessary to prechill pipette tips, tubes, plates, or other objects that may come in contact with the extract. Since smaller volumes warm more quickly, partial tubes and aliquots should be kept on ice to prevent premature gelling.

Thawing of Geltrex, dilution in DMEM/F12 and aliquotting fo..

1

🔥 4 °C

Thaw one tube of Geltrex (1-5 mL) slowly at 2-8°C.

Prepare 20X 1.5 ml Eppendorfs for aliquotting of Geltrex, label: 50X

Mix Geltrex by slowly pipetting solution up and down; be careful not to introduce air bubbles.

On ice: in a 15 ml Falcon tube, mix Geltrex 1:1 with **cold** DMEM/F12 (Sigma #D6421) and aliquot at 0.5 ml.

Store the aliquots at -20C.

(REMEMBER TO CHECK EXPIRY DATE ON EVERYTHING)

Plate coating

2 Take plates that require coating and label.

Thaw an aliquot of Geltrex/DMEM/F12 (1:1 mix) on ice.

Prepare an aliquot of 22 ml cold DMEM/F12 = keep on ice. 🔥 4 °C

On ice: Dilute 450 ul of 1:1 Geltrex (50X) ~1:50 into 22 ml **cold** DMEM/F12 medium for a final dilution of 1:100. Empirical determination of the optimal coating concentration for your application may be required. Volumes can be adjusted accordingly.

Add a sufficient amount of diluted Geltrex solution to cover the entire area onto growth surface (100 ul/well in 96-well plate; 0.5 ml/well for 24-well and 4-well Ibidi imaging dish; 1 ml for 35 mm dish, 3 ml for 60 mm dish, i.e. 0.14 ml/cm² --> 1 ml per 6-well dish; 8 ml per T75 flask).

Coat the dish and place in incubator (37C) for 60 min (no need to seal). 🔥 37 °C

Afterwards, take the plates that you will need for cell seeding out to room temperature for minimum 30 min. Store the remaining plates at 4C for up to 2 weeks (best for only 1 week to be on the safe side; parafilm to avoid drying out).

Do not allow coated surface to dry out and maintain a storage temperature of 2 to 8°C to avoid premature gelling. 🔥 4 °C

At time of use, aspirate Geltrex™ coating and immediately plate cells in pre-equilibrated cell

culture medium. No rinsing!