

Oct 17, 2019

## -Colony Formation Unit Assay

DOI

[dx.doi.org/10.17504/protocols.io.8bkhskw](https://dx.doi.org/10.17504/protocols.io.8bkhskw)

NUS iGEM<sup>1</sup>

<sup>1</sup>National University of Singapore



OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.8bkhskw](https://dx.doi.org/10.17504/protocols.io.8bkhskw)

**Protocol Citation:** NUS iGEM 2019. Colony Formation Unit Assay. [protocols.io](#)

<https://dx.doi.org/10.17504/protocols.io.8bkhskw>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

We use this protocol and it's working

**Created:** October 17, 2019

**Last Modified:** September 12, 2023

**Protocol Integer ID:** 28748

## Materials

MATERIALS



- 1 Dilute cell culture with 1x PBS to a final concentration of  $10^6$ ,  $10^7$  and  $10^8$
  
- 2 Aliquot  40  $\mu\text{L}$  of each sample onto the agar plate containing appropriate antibiotic

## Protocol



NAME

### Preparation of LB Agar

CREATED BY

NUS iGEM

[PREVIEW](#)

- 2.1 Weigh  32 g of LB agar powder.

 LB Agar, powder (Lennox L agar) Thermo Fisher Catalog #22700025

- 2.2 Add the powder into  1 L of deionized water.

 Water refers to sterilized deionized water

- 2.3 Autoclave entire bottle of LB media.

- 3 Spread the cells evenly

- 4 Incubate at  37 °C overnight