SEP 20, 2019

Colony PCR

宏亮 董¹

¹Northeast Forest University

2019 iGEM NEFU_China

Tech. support email: shengyiyanwork@gmail.com

宏亮 董

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7h2hj8e

Protocol Citation: 宏亮 董
2019. Colony PCR.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.7h2hj8e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 20, 2019

Last Modified: Sep 20, 2019

PROTOCOL integer ID:
27930

Keywords: Colony PCR

ABSTRACT

This PCR method can be used to screen for inserted target genes or DNA sequencing analysis.

MATERIALS

2×high Taq Master Mix (Enzyme)

Template

F/R primers

ddH₂O

Bio-rad S1000TM Thermo Cycler.

SAFETY WARNINGS

Please wear gloves for the experiment, don't try to touch the lid after PCR program initiation.

BEFORE START INSTRUCTIONS

Synthesize primers in advance.

- 1 Pick colonies as the template for colony PCR. The number picked for each plate depends on the difference between the positive and negative controls.

2×high Taq Master Mix (Enzyme)	5 µl
Template	0.4 µl
Forward Primer (10 µM)	0.4µl
Reverse Primer (10 µM)	0.4 µl
ddH2O	3.8 µl

Fill the rest with water.

- 2 Test digest performed and products analysed using agarose gel electrophoresis to confirm if correct construct was present.

- 3 Use colony PCR to enlarge colony numbers. Then only the positive clones were mini-prepped.