MAY 07, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.zp7f5rn

External link:
<https://doi.org/10.1371/journal.pone.0216363>

Protocol Citation: Barbara Rizzacasa, Elena Morini, Ruggiero Mango, Chiara Vancheri, Simone Budassi, Gianluca Massaro, Sara Maletta, Massimiliano Macrini, Silvio D'Annibale, Francesco Romeo, Giuseppe Novelli, Francesca Amati 2019. Comparative Ct method quantification (2- Δ Ct method).. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.zp7f5rn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Apr 02, 2019

Last Modified: May 07, 2019

🌐 Comparative Ct method quantification (2- Δ Ct method).

📁 PLOS One

Barbara Rizzacasa¹, Elena Morini¹, Ruggiero Mango², Chiara Vancheri¹, Simone Budassi², Gianluca Massaro², Sara Maletta¹, Massimiliano Macrini², Silvio D'Annibale³, Francesco Romeo^{4,5}, Giuseppe Novelli¹, Francesca Amati^{6,7}

¹[Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, Italy];

²[Complex Operative Unit of Cardiology (UOC), Policlinic Tor Vergata, Rome, Italy];

³[Department of System Medicine, University of Rome Tor Vergata, Rome, Italy];

⁴[Complex Operative Unit of Cardiology (UOC), Policlinic Tor Vergata, Rome, Italy];

⁵Department of System Medicine, University of Rome Tor Vergata, Rome, Italy];

⁶[Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, Italy];

⁷Department of Human Sciences and Quality of Life Promotion, University San Raffaele, Rome, Italy]

Francesca Amati

ABSTRACT

For the evaluation of the expression level of each miRNA was used the Ct (threshold cycle) method quantification. The threshold cycle (Ct) is defined as the PCR cycle at which the fluorescent signal of the reporter dye crosses an arbitrarily placed threshold. This method allows to quantify the absolute expression of each miRNAs in each sample analyzed and then to calculate the different expression of each miRNA in sample versus the controls.

Keywords: qRT-PCR,
threshold cycle, microRNAs
quantification

Evaluation of the absolute expression of miRNAs in each sa..

- 1 For each sample is obtained an amplification curve in which the Ct is inversely proportional to the initial sample quantity;
- 2 Normalize the data to an internal control and calculate $DCt = (Ct \text{ gene of interest} - Ct \text{ internal control})$;
- 3 Calculate $2^{-\Delta Ct}$ value that is the measure of the microRNA expression level in each sample analyzed.

Evaluation of the expression of each miRNA in two differen...

- 4 Calculate $DDCt = [(Ct \text{ gene of interest} - Ct \text{ internal control}) \text{ sample A}] - [(Ct \text{ gene of interest} - Ct \text{ internal control}) \text{ sample B}]$;
- 5 Calculate $2^{-\Delta\Delta Ct}$ value that is the measure of the microRNA expression in each sample analyzed compared with control sample.