SEP 10, 2018

Competition among *Aedes aegypti* larvae in microcosms

PLOS One

Kurt Steinwascher¹

¹retired

Kurt Steinwascher
retired

ABSTRACT

Female larvae of the *Aedes aegypti* mosquito outcompete male larvae for food. Male larvae pupate earlier and at smaller masses in vials where competition is most intense. Competition at the higher densities causes the food/larva to appear lower than the equivalent mg/larva at lower densities for both sexes. The total food per vial appears to change the nature of the exploitative competition among females, which affects the competition among males. No interference competition was observed; females may dominate the competition by their larger size (increased filtering ability) and by retaining food particles within their guts to extract a larger fraction of their nutrients. Pupal mass for both sexes is primarily affected by food/larva; density affects the pupal mass through competition and the total food/vial (the interaction between food and density) and this interaction is different for the two sexes. Age at pupation is more affected by density than by food/larva and the interaction between food and density also differs across the two sexes for this variable. The nature of competition between and within the sexes at various food levels and densities is described. The implications of this complex response to food level and density among larvae on the ecology of this species is considered.

GUIDELINES

It is easier to create a standardized solution of baker's yeast and dilute it down to the concentration necessary to add the calculated total mg per vial for each treatment. However, the yeast must be completely dissolved in the solution and it must be repeatedly stirred to keep it from clumping or settling.

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.tcweixe](https://doi.org/10.17504/protocols.io.tcweixe)

External link:
<https://doi.org/10.1371/journal.pone.0202455>

Protocol Citation: Kurt Steinwascher 2018. Competition among *Aedes aegypti* larvae in microcosms. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.tcweixe>

MANUSCRIPT CITATION: Steinwascher K (2018) Competition among *Aedes aegypti* larvae. PLoS ONE 13(11): e0202455. doi: [10.1371/journal.pone.0202455](https://doi.org/10.1371/journal.pone.0202455)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 09, 2018

PROTOCOL integer ID:
15478

Keywords: *Aedes aegypti*,
age at pupation, ANOVA,
competition, exploitative,
female advantage, filter
feeding, food x density,
growth rate, interference,
larval density, larval
development, life history,
MANOVA, multivariate analysis
of variance, pupal mass,
pupation, sex ratio, size
distribution, survival

- 1 For experiment 1: Obtain mosquito eggs from an *A. aegypti* colony after feeding females on a mouse.
- 2 Hatch the eggs in distilled water, immersing them for at least 2 hours.
- 3 Assign random numbers to treatments: 20 treatments (4 food levels crossed with 5 densities, see below) with 5 replicates each.
- 4 Number 100 flat-bottomed shell vials (25 mm diameter x 95 mm tall) with random number and treatment.
- 5 Fill shell vials with 20 ml distilled water in each.
- 6 Weigh out baker's yeast and dissolve it in distilled water to achieve the following concentrations: 2 mg/larva, 3 mg/larva, 4 mg/larva and 5 mg/larva crossed with 5 densities (4, 5, 6, 7 and 8 larvae per vial). [2 mg/larva with 4 larva per vial requires a total amount of yeast of 8 mg, dissolved in 0.5 ml distilled water and added to the 20 ml in the appropriate vials—5 replicates of each

treatment, for 20 treatments.]

- 7 Count 4, 5, 6, 7 or 8 larvae into each vial according to the treatment marked on the vial.
- 8 Arrange vials by number (randomized sequence) and cover with cheese cloth to prevent food or other mosquitoes from entering the vials.
- 9 Store vials in a room at ambient temperatures (in Florida) with a temperature range of 18 degrees C to 33 degrees C.
- 10 Day zero is the day the experiment begins. Check the vials for pupae beginning on day 4 (this is the day before any larvae are expected to pupate).
- 11 Remove each pupa into a holding shell vial with 10 ml distilled water. Record the original vial number, the treatment, the date (day of pupation). To pick up a pupa without disturbing the treatment vial, use a fire-polished wide mouth glass pipette (about 2 mm diameter) to capture the pupa at the water surface and transfer it to the holding vial. Capillary forces will capture the pupa, but you may have to gently blow the pupa out of the tube into the holding vial. Put only 1 pupa in each holding vial.
- 12 Remove each pupa from the holding vial by pipette as described, blot it on a piece of paper towel, pick it up gently with fine-tipped forceps and place it on a balance. Weigh it to the nearest 0.01 mg. Record the weight. Transfer it to the stage of a 10 x stereoscopic microscope and determine the sex of the pupa. Record the sex. Transfer it back to a vial of water to complete pupation (and eventually to rejoin the mosquito colony).
- 13 Continue until all larvae have pupated or died.
- 14 For experiment 2: Obtain mosquito eggs from an *A. aegypti* colony after feeding females on a mouse.

- 15 Hatch the eggs in distilled water, immersing them for at least 2 hours.
- 16 Assign random numbers to treatments: 4 treatments (2 food levels crossed with 2 densities, see below) with 40 replicates each.
- 17 Number 160 flat-bottomed shell vials (25 mm diameter x 95 mm tall) with random number and treatment.
- 18 Fill shell vials with 20 ml distilled water in each.
- 19 Weigh out baker's yeast and dissolve in distilled water to achieve the following concentrations: 3 mg/larva and 4 mg/larva crossed with 2 densities (5 and 6 larvae per vial). [3 mg/larva with 5 larva per vial requires a total amount of yeast of 15 mg, dissolved in 0.5 ml distilled water and added to the 20 ml in the appropriate vials—40 replicates of each treatment, for 4 treatments.]
- 20 Count 5 or 6 larvae into each vial according to the treatment marked on the vial.
- 21 Arrange vials by number (randomized sequence) and cover with cheese cloth to prevent food or other mosquitoes from entering the vials.
- 22 Store vials in an insectary at 26 degrees C and 12/12 light/dark cycle for the first four days and overnight thereafter.

- 23** Day zero is the day the experiment begins. Check the vials for pupae beginning on day 4 (this is the day before any larvae are expected to pupate).
- 24** Remove each pupa into a holding shell vial with 10 ml distilled water. Record the original vial number, the treatment, the date (day of pupation). To pick up a pupa without disturbing the treatment vial, use a fire-polished wide mouth glass pipette (about 2 mm diameter) to capture the pupa at the water surface and transfer it to the holding vial. Capillary forces will capture the pupa, but you may have to gently blow the pupa out of the tube into the holding vial. Put only 1 pupa in each holding vial.
- 25** Remove each pupa from the holding vial by pipette as described, blot it on a piece of paper towel, pick it up gently with fine-tipped forceps and place it on a balance. Weigh it to the nearest 0.01 mg. Record the weight. Transfer it to the stage of a 10 x stereoscopic microscope and determine the sex of the pupa. Record the sex. Transfer it back to a vial of water to complete pupation (and eventually to rejoin the mosquito colony).
- 26** Continue until all larvae have pupated or died.