AUG 11, 2020

Competitive ELISA to study the inhibition of the SpA-binding to Ab-3 by Ab-2.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjkbkksn

Protocol Citation: Angel A Justiz-Vaillant 2020. Competitive ELISA to study the inhibition of the SpA-binding to Ab-3 by Ab-2.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjkbkksn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 11, 2020

Last Modified: Aug 11, 2020

PROTOCOL integer ID:
40291

ABSTRACT

This immunoassay tested the ability of anti-anti-SpA (Ab-2) to inhibit or interfere with the binding of anti-anti-anti-SpA (Ab-3) to the original antigen- staphylococcal protein A (SpA). It is a confirmation test of the functional capacity of Ab-2. In this ELISA, both Ab-2 and the antigen SpA compete for binding to Ab-3. When Ab-2 is present, it blocks the binding of HRP-labeled SpA to Ab-3, resulting in inhibition of reaction color development.

GUIDELINES

Use appropriate positive and negative controls, as well as blanks.

MATERIALS

MATERIALS

- ⊗ Substrate (TMB) **Millipore Sigma Catalog #EZMADP-60K(kit)**
- ⊗ Stop Solution (0.3M HCl) **Millipore Sigma Catalog #EZMADP-60K(kit)**
- ⊗ 96-Well Microtiter™ Microplates, Polystyrene, 280µL, Nonsterile, V-bottom **Thermo Fisher Catalog #2605**
- ⊗ Pierce™ Recombinant Protein A, Peroxidase Conjugated **Thermo Fisher Catalog #32400**

SAFETY WARNINGS

! Do not use any reagent where damages to the packaging has occurred.

- 1 Prepare materials and ELISA buffer solutions and reagents.

- 2 Pipette 53 μ l of purified Ab-3 (100 μ g/ml) mixed with 2.8 ml of coating buffer into each well.
- 3 Incubate the microplate at 37°C for 4 hrs.
- 4 Aspirate the contents of the wells.
- 5 Fill each well with an appropriately diluted washing solution and aspirate.
- 6 Wash the microplate 3 times.
- 7 Pipette 53 μ l of serial dilutions of pooled Ab-2 (100 μ g/ml) in triplicates.
- 8 Incubate the microplate at RT for 1 hr.
- 9 Rewash the microplate filling each well with 100 μ l of washing buffer.

- 10** Pipette 53 μ l of commercially available peroxidase-labeled SpA conjugate (Sigma-Aldrich) diluted 1:5000 to each well.
- 11** Reincubate the microplate at RT for 1 hr.
- 12** Repeat step 5.
- 13** Pipette 53 μ l of TMB (Sigma-Aldrich) to each well.
- 14** Incubate in the dark at RT for 14 min.
- 15** Measure absorbance at 450 nm in a microplate reader and analyze the results