AUG 27, 2019

CPEC Protocol

In 1 collection

N.J. Hillson^{1,2}

¹JBEI; ²LBNL

Michael J Fero

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.6t7hern

Protocol Citation: N.J. Hillson
2019. CPEC Protocol.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.6t7hern>

MANUSCRIPT CITATION:

J. Quan and J. Tian, "Circular polymerase extension cloning of complex gene libraries and pathways," PloS one, vol. 4, no. 7, p. 6441, 2009.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use these protocols in our group and they work.

Created: Aug 26, 2019

Last Modified: Aug 27, 2019

PROTOCOL integer ID:

27231

Measure the DNA concentration (ng/ml) of each assembly piece

- 1 measure the DNA concentration (ng/ml) of each assembly piece.

- 2 Add 100 ng of the linearized vector backbone and equimolar amounts of the other assembly pieces to a 25 ml total volume assembly reaction mixture as follows:

linearized vector backbone (100 ng)
+ each additional assembly piece (to equimolar with backbone)
+ 5 ml 5X HF Phusion Reaction Buffer
+ 1 ml 10 mM dNTPs
+ 0.75 ml DMSO
+ 0.5 ml 2U/ml Phusion Polymerase
+ _____dH₂O to
25 ml

- 3 Perform the assembly reaction in a thermocycler as follows:

30 sec @ 98 C 1 cycle
10 sec @ 98 C }
30 sec @ 55 C } 1 to 15 cycle(s)**
length* (kb) x 15 sec @ 72 C }
10 min @ 72 C 1 cycle

*The total length of the assembled product (in kb)

**The number of repeated cycles should exceed the number of assembly pieces

Transform 5 ml of the assembly reaction into 100 ml of competent E. coli and/or run a diagnostic agarose gel to check for successful assembly.