JUN 10, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.quxdwxn

Protocol Citation: Bulent Arman Aksoy, Pinar Aksoy, Jeff Hammerbacher 2018. CRISPR/Cas9-based knock-out in human primary T cells (24-well setup). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.quxdwxn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 10, 2018

Last Modified: Jun 10, 2018

PROTOCOL integer ID:
 12919

🌐 CRISPR/Cas9-based knock-out in human primary T cells (24-well setup)

📁 In 6 collections

Bulent Arman Aksoy¹, Pinar Aksoy¹, Jeff Hammerbacher¹

¹Medical University of South Carolina

Hammer Lab

Tech. support phone: +18437924527 email: arman@hammerlab.org

Bulent Arman Aksoy

Medical University of South Carolina

ABSTRACT

This protocol has been optimized for [Thermo Fisher's Neon Electroporation System](#) and is based on the report [Reprogramming human T cell function and specificity with non-viral genome targeting](#). We have used [TrueCut™ Cas9 Protein v2](#) and reached very comparable KO efficiencies to those previously shown by others.

GUIDELINES

To reduce variation per electroporation sample and to account for practical issues of electroporation (e.g. sparks), we have also found that a 24-well plate setup where we seeded cells from three independent electroporation reactions (within 10-ul tip) into a single well with 500 ul to 1 ml culture media in it had been the most feasible approach. This setup is especially helpful during initial optimization experiments where the goal is to screen many conditions. Pooling three independent reactions into a single sample provides a virtual buffer against variation across electroporation samples and yields enough treated cells for common assaying approaches, such as Western Blotting. We always recommend optimizing or screening using 10-ul tip setup and then scaling the reactions up to 100-ul once the optimal settings are known.

MATERIALS

MATERIALS

⊗ TrueCut™ Cas9 Protein v2 **Thermo Fisher Scientific Catalog #A36497**

⊗ TrueGuide™ tracrRNA **Thermo Fisher Scientific Catalog #A35506**

⊗ Neon™ Transfection System 10 µL Kit **Thermo Fisher Scientific Catalog #MPK1096**

Cas9 RNP preparation

- 1 RNase Zap everything before you start
- 2 Resuspend the crRNA (2 nmol) in 100 ul (making 20 uM) and tracrRNA (20 nmol) in 1 ml RNA storage buffer (making 20 uM). Aliquot and keep at -80°C.
- 3 Need 0.375ul crRNA and 0.375 ul tracrRNA per well (this way, we'll use 7.5pmol sgRNA per 200K cells as Neon protocol suggests).
- 4 Mix 1.5 ul crRNA and 1.5 ul tracrRNA (for 4 rxns) in a PCR tube.
- 5 Keep the RNA mix at 95°C for 5 minutes
00:05:00
- 6 Then keep the RNA mix at at 37°C for 25 mins.
00:25:00
- 7 For Cas9, Neon suggests 1250 ng Cas9 protein per 200K cells. Our Cas9 is at 5 mg/ml concentration. So, we will need 0.25 ul Cas9/200K cells.
- 8 After incubation of the sgRNA, slowly add 1ul Cas9 (for 4 reactions) in the PCR tube, mix and incubate at 37°C for 15 mins.
00:15:00

Cell preparation and electroporation

- 9 Need 200K cells for one electroporation event. The cells should be in 9 ul T buffer (9 ul T buffer reaction), so that the total volume (9 ul cells + 1 ul RNP mix) will be 10ul for the Neon 10 tip.
- 10 Debead and count the activated cells.
- 11 We need $200,000 \times 3 \times 24$ (14.4 million cells).
- 12 To be safe, assume 4 reactions well so we will need $200,000 \times 4 \times 24$ (19 million) cells.
- 13 19 million cells are actually good for 96 wells.
- 14 So, we need $96 \times 9 = 864$ ul T buffer
- 15 After debeading and counting the cells, spin them down at 200 x g for 7 mins.
- 16 Aspirate the media as much as possible.
- 17 Resuspend the pellet in 864 ul T buffer.

18 Add 36 ul of cell mix to each PCR tube and mix well.

19 The Cas9 RNP and cells are ready for electroporation

20 Electroporation at 1600 V 10 ms 3 pulses

Equipment

Thermo Fisher Scientific Neon™ Transfection System

MPK5000S

NAME

BRAND

SKU

21 Seed the electroporated cells on the prepared 24-well-plate with warm T cell media.

Profiling

22 The time to profile CRISPR/Cas9 treated cell depends on the particular assay of interest but in general, cells can be profiled via flow cytometer for surface proteins and via Western Blot for internal proteins 3 days after electroporation.