


Dec 12, 2019

## CRISPR/Cas9 based knockout generation in *Aurantiochytrium limacinum* (ATCC MYA-1381)

DOI

[dx.doi.org/10.17504/protocols.io.baeyibfw](https://dx.doi.org/10.17504/protocols.io.baeyibfw)



Anbarasu Karthikaichamy<sup>1</sup>, Jackie Collier<sup>2</sup>

<sup>1</sup>University of Illinois at Urbana-Champaign; <sup>2</sup>State University of New York at Stony Brook

Collier Lab



Anbarasu Karthikaichamy

University of Illinois at Urbana-Champaign

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.baeyibfw](https://dx.doi.org/10.17504/protocols.io.baeyibfw)

**Protocol Citation:** Anbarasu Karthikaichamy, Jackie Collier 2019. CRISPR/Cas9 based knockout generation in *Aurantiochytrium limacinum* (ATCC MYA-1381). [protocols.io https://dx.doi.org/10.17504/protocols.io.baeyibfw](https://dx.doi.org/10.17504/protocols.io.baeyibfw)

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

We use this protocol and it's working

**Created:** December 12, 2019

**Last Modified:** December 12, 2019

**Protocol Integer ID:** 30904

**Keywords:** Stramenopile, Thraustochytrid, *Aurantiochytrium*, CRISPR, Protists, Electroporation

## Abstract

Transformation protocol for electroporation of CRISPR/Cas9 ribonucleoprotein (RNP) complex in urantiochytrium limacinum (ATCC MYA-1381; Stramenopile/ Heterokont, Thraustochytrid).

## Materials

### MATERIALS

 NEPA21 Super Electroporator **NEPAGENE**

 EnGen® sgRNA Synthesis Kit S. pyogenes **New England Biolabs Catalog #E3322S**

 Cas9 Nuclease S. pyogenes **New England Biolabs Catalog #M0386S**

 Monarch® RNA Cleanup Kit **New England Biolabs Catalog #T2050S**

### STEP MATERIALS

 EnGen® sgRNA Synthesis Kit S. pyogenes **New England Biolabs Catalog #E3322S**

 Cas9 Nuclease S. pyogenes **New England Biolabs Catalog #M0386S**

 Monarch® RNA Cleanup Kit **New England Biolabs Catalog #T2050S**

## Protocol materials

 EnGen® sgRNA Synthesis Kit S. pyogenes **New England Biolabs Catalog #E3322S**

 Cas9 Nuclease S. pyogenes **New England Biolabs Catalog #M0386S**

 Monarch® RNA Cleanup Kit **New England Biolabs Catalog #T2050S**

 NEPA21 Super Electroporator **NEPAGENE**

 EnGen® sgRNA Synthesis Kit S. pyogenes **New England Biolabs Catalog #E3322S**

 Cas9 Nuclease S. pyogenes **New England Biolabs Catalog #M0386S**

 Monarch® RNA Cleanup Kit **New England Biolabs Catalog #T2050S**

 EnGen® sgRNA Synthesis Kit S. pyogenes **New England Biolabs Catalog #E3322S**

 Cas9 Nuclease S. pyogenes **New England Biolabs Catalog #M0386S**

 Monarch® RNA Cleanup Kit **New England Biolabs Catalog #T2050S**

## Designing oligos for gRNA synthesis

40m

- 1 Use ChopChop tool ([chopchop.rc.fas.harvard.edu](http://chopchop.rc.fas.harvard.edu)) to select the gRNA of interest. *Aurantiochytrium limacinum* genome is available on ChopChop, and the gene of interest can be accessed using the gene co-ordinates of the JGI assembly <https://mycocosm.jgi.doe.gov/Aurli1/Aurli1.home.html>(For example: scaffold\_10:1222284-1223740 (+)). 10m
- 2 The output will list the gRNAs in a ranked order, considering the self-complementarity of the gRNA, efficiency, and off-target effects. Select the highly ranked gRNA and proceed for oligo designing for in vitro transcription. 10m
- 3

### Note

Design target specific oligos following the [NEB protocol](#)

Select 20 nt gRNA sequences using the ChopChop web tool (do not include the PAM sequences), and add 'G' to the 5' end of the sequence, only if there is no 'G' at the 5' end.

- 4 To the 5' end; append T7 promoter sequence: **TTCTAATACGACTCACTATA**  
To the 3' end; append 14 nucleotide overlap sequence: **GTTTTAGAGCTAGA**  
The final oligo sequence should look like 5'  
**TTCTAATACGACTCACTATAG(N)<sub>20</sub>GTTTTAGAGCTAGA** 3', (N)<sub>20</sub> is the gRNA sequence.  
This is the sequence of the oligo to be ordered.

10m

### Note

**IMPORTANT:** While the oligos are shipped, we recommend starting the pre-culture (step8) at this stage since it takes 2-3 days to grow.

## gRNA synthesis and purification

1h 30m

- 5

1h



### Note

Wear gloves and use nuclease-free tubes and reagents. Reactions should be assembled in microfuge tubes or PCR strip tubes.

Perform gRNA synthesis following the [EnGen®sgRNA Synthesis Kit](#).

 EnGen® sgRNA Synthesis Kit S. pyogenes New England Biolabs Catalog #E3322S

- 6 Purify the synthesized gRNA using Monarch RNA Cleanup Kit (50 µg)

30m

 Monarch® RNA Cleanup Kit New England Biolabs Catalog #T2050S

Quantify the gRNA using UV-Vis Nano-spectrometer.

Label and store the gRNA at  -20 °C for short term storage and at  -80 °C for long term storage.

## Ribonucleoprotein (RNP) preparation

30m

- 7 Thaw the gRNA and 1x PBS on ice. Spin down all the reagents before using.

30m





For a total reaction volume of 5µl,

1. Add  120 pmol (final) gRNA and  104 pmol (final) Cas9 protein.
2. Adjust the reaction volume to 5µl using 1x PBS.

 Cas9 Nuclease S. pyogenes New England Biolabs Catalog #M0386S

Mix the reagents by gently pipetting up and down. After a brief spin, incubate the tube at

 Room temperature for  00:20:00 .

Label and store the RNP (Cas9/gRNA) complex in  -20 °C until further use.

### Note

#### Optional

[Check the efficiency of RNP by performing an in vitro cleavage assay.](#)

## Grow cells

4d

- 8 1. Start a pre-culture  96:00:00 (4 days) prior to electroporation by inoculating  5 mL of GPY (0.5% Yeast Extract, 1% Peptone, 3% D+-Glucose, 1.8% instant ocean) in  15 mL tube with a colony of *Aurantiochytrium limacinum* (ATCC MYA-1381). Incubate  Overnight at  28 °C .
2. Use preculture to inoculate  46 mL of GPY in  250 mL flask. Culture for 3 days at  28 °C ,  171 rpm .

## Prepare reagents for Electroporation

2h

- 9 1. Prepare GPy's media (3% glucose, 0.6% peptone, 0.2% yeast extract,  50 millimolar (mM) sucrose, 1.8% instant ocean)
2. Prepare GPy's ampicilin (100µg/ml) plates (2%agar)
3. Prepare GPy's zeocin (100µg/ml) and ampicilin (100µg/ml) plates (2%agar)

### Note

GPy's zeocin + ampicillin plates are used if homology directed repair (HDR) template with Zeo<sup>R</sup> is used.

1. Prepare 1X BSS ( 10 millimolar (mM) KCl,  10 millimolar (mM) NaCl, and  3 millimolar (mM) CaCl2)
2. Prepare  50 millimolar (mM) sucrose solution
3. Sterilize by autoclave or filter sterilization as appropriate.

### CITATION

Mariana Rius, Jackie Collier. Electroporation of *Aurantiochytrium limacinum* (ATCC MYA-1381).

LINK

[dx.doi.org/10.17504/protocols.io.qjcduiw](https://dx.doi.org/10.17504/protocols.io.qjcduiw)

## Prepare cells for electroporation

45m

- 10 1. Count cells using haemocytometer. Cell density should be around  $\sim 5 \times 10^7$  cells/ml.  
2. Add 1.5 ml of cells to a microcentrifuge tube for each electroporation reaction.  
Centrifuge  
3.  11000 rpm, 4°C, 00:05:00  
4. Decant the supernatant. Add 500  $\mu$ L chilled 1X BSS (  
[M] 10 millimolar (mM) KCl,  
[M] 10 millimolar (mM) NaCl, and  
[M] 3 millimolar (mM) CaCl<sub>2</sub>). Centrifuge  
 11000 rpm, 4°C, 00:05:00  
5. Decant the supernatant. Add  500  $\mu$ L chilled [M] 50 millimolar (mM) sucrose.  
Centrifuge  11000 rpm, 4°C, 00:05:00 . Repeat thrice (3x).  
6. Re-suspend by scraping cell mass off side of tube and vortex.

## Add RNP

5m

- 11 1. Add  5  $\mu$ L of RNP complex mixture to suspended cells.  
2. Incubate on ice  00:05:00 .  
3. Transfer to chilled electroporation cuvette (0.2cm; BioRad).  
4. Keep the cuvettes on ice.  
5. Keep appropriate negative control using the same volume of elution buffer or water.

## Electroporation

10m

12



## Equipment

NEPA21 Super Electroporator

NAME

NEPAGENE

BRAND

NEPA21

SKU

[http://www.nepagene.jp/e\\_products\\_nepagene\\_0001.html](http://www.nepagene.jp/e_products_nepagene_0001.html)<sup>LINK</sup>



Clean the loading pedestal and the cuvette before loading.

Set the parameters for poring and transfer pulse,

### Poring pulse:

Voltage (V)	Pulse length (ms)	Length interval (ms)	No. of pulse	Decay rate (%)	Polarity
275	8	50	2	10	+

### Transfer pulse:

Voltage (V)	Pulse length (ms)	Length interval (ms)	No. of pulse	Decay rate (%)	Polarity
20	50	50	1	40	+/-

The capacitance and resistance is set at 125  $\mu\text{F}$ , and 1000  $\Omega$  respectively.

Pulse the cells and record the time constant. Take out the cuvette and place it on ice.

Repeat the steps for the rest of the samples.

## Outgrowth and plating

1h 30m

- 13 1. Add 1ml GPYs media to the cuvette and carefully aspirate out the cells on to a fresh micro centrifuge tube.
2. Label the corresponding tubes and Incubate at  28 °C for  01:00:00 without shaking.
3. Centrifuge the cells at  11000 rpm, 25°C, 00:05:00, and discard the supernatant.
4. Re-suspend the cells in remaining supernatant by pipetting up and down and also by vortexing.
5. Dilute the cells appropriately (100x, 1000x) in  100 µL of fresh GPYs media.
6. Plate the cells on appropriate GPYs plates, we usually include ampicillin to control bacterial contamination, and incubate at  28 °C.

1h 30m

## Monitor transformant colonies

2d

- 14 Colonies of *Aurantiochytrium limacinum* transformants will appear 2 days after plating. Streak individual colonies on fresh GPYs plate, and observe the phenotype/genotype. Include wild type (WT) for comparison.

### Note

The transformant selection/screening can vary depending on the target gene and the selection marker used. In our case, we knocked-out carotenogenic gene, and the phenotype selection was based on colony colour.

## Citations

### Step 9

Mariana Rius, Jackie Collier. Electroporation of *Aurantiochytrium limacinum* (ATCC MYA-1381)  
[dx.doi.org/10.17504/protocols.io.qjcduiw](https://dx.doi.org/10.17504/protocols.io.qjcduiw)