

Nov 12, 2023

Version 1

CRISPR knock-in (endogenous tagging) V.1

DOI

<https://dx.doi.org/10.17504/protocols.io.e6nvwdmqwlmk/v1>

Leonardo Parra-Rivas¹

¹University of California, San Diego



Leonardo A Parra-Rivas

University of California, San Diego

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.e6nvwdmqwlmk/v1>

Protocol Citation: Leonardo Parra-Rivas 2023. CRISPR knock-in (endogenous tagging). protocols.io

<https://dx.doi.org/10.17504/protocols.io.e6nvwdmqwlmk/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: November 12, 2023



Last Modified: November 12, 2023

Protocol Integer ID: 90814

Keywords: crispr knock, crispr, endogenous tagging

Abstract

CRISPR knock-in (endogenous tagging)

Plasmid construction

1

For the C-terminus knock-in experiment, the sgRNAs and the homology-independent donor templates were generated following strategies similar to those described previously

Citation

Ogawa Y, Rasband MN (2021)
. Endogenously expressed Ranbp2 is not at the axon initial segment.

<https://doi.org/10.1242/jcs.256180>

LINK

Briefly, the coding sequence of *SNCA1* –the gene encoding α -syn at exon 6 was targeted by a sgRNA, and the oScarlet tag (a variant of mScarlet that is engineered to reduce aggregation; a gift from Karl Deisseroth, Addgene plasmid #137135) was knocked in using a homology-independent mechanism .

- 2 The donor sequence was designed following the SATI knock-in vector with slight modifications for the intron knock-in experiment. Briefly, the following sequences were directly conjugated as a donor tag; *SNCA1* intron 4 (only the sequence after the gRNA targeting site), *SNCA1* exon 5 (with wild type or mutation sequences; S129A or S129D), *SNCA1* exon 6 (coding site until stop codon), 3X GGGGS linker, oScarlet (without a start codon and with a stop codon at the end), and *SNCA1* 3' UTR. The *SNCA1* intron 4 was targeted by a sgRNA (TTCTAAGTGATACCAAACCAC), and the donor tag was homology-independently knocked in. The plasmid PX552 (RRID:Addgene_60958) (a gift from Feng Zhang) was digested with a NotI restriction enzyme (New England Biolabs, Cat#R3189L) and used as a plasmid backbone.

- 3 Oligonucleotides were ordered from Sigma. The genomic sequence of *SNCA1* was partially synthesized by Twist Bioscience, and for the rest, it was PCR amplified from a purified genome of a C57BL/6J mouse. DNA fragments were ligated together using the In-Fusion Snap Assembly Master Mix (Takara, Cat# 638947) or DNA Ligation Kit Mighty Mix (Takara, Cat#6023). The sgRNA sequence for knock-in is listed in the supplemental materials. The AAV-SpCas9 plasmid PX551 (RRID:Addgene_60957 (a gift from Feng Zhang)) was modified by removing the HA tag.

Adeno-associated virus (AAV) production, neuronal transduction, and quantification of synapses

- 4

Small-scale AAV cell lysates were produced using the AAVpro Purification Kit (All Serotypes) (Takara, Cat# 6666) with slight modifications. Briefly, HEK293T cells (RRID:CVCL_0063) were triple transfected with the AAV targeting plasmid, helper plasmid (Agilent Technologies, Cat # 240071), and serotype PHP.S plasmid pUCmini-iCAP-PHP.eB (RRID:Addgene_103005)(a gift from Viviana Gradinaru) with PEI Max (Polysciences, Cat#24765).
- 5 The medium was changed the next day of transfection, and cells were incubated for 3 days after transfection. HEK cells were then collected and lysed with the AAV Extraction Solution A plus.
- 6 The extracted solution was centrifuged at 10,000 x g for 10 min to remove debris and mixed with Extraction Solution B. This small-scale AAV solution was stored at -80 until use. For viral transduction, hippocampal neurons were plated at 60,000 cells/cm² density and infected 4 hours later with 10ul of the AAVs expressing SpCas9 and sgRNAs/donor.
- 7 The media was replaced 2 days after infection. Before imaging or genomic analysis, the transduced neurons were cultured to maturity (DIV-17-DIV-21).



- 8 For quantification of the knock-in α -syn:o-Scarlet fluorescence at synapses, images were first background-corrected, small ROIs were manually placed over ~20-30 synapses on each image, and average intensities were calculated – all using dropdown menus in MetaMorph Microscopy Automation and Image Analysis Software (RRID:SCR_002368) (<https://www.moleculardevices.com/products/cellular-imaging-systems/acquisition-and-analysis-software/metamorph-microscopy#gref>). The resulting datasets were statistically analyzed using GraphPad Prism [(RRID:SCR_002798) <http://www.graphpad.com/>].

Citations

Step 1

Ogawa Y, Rasband MN. Endogenously expressed Ranbp2 is not at the axon initial segment. <https://doi.org/10.1242/jcs.256180>