

Jul 10, 2019

Version 1

🌐 Cultivating Melanized Fungi from Biological Soil Crust and Rock Surfaces V.1

DOI


[dx.doi.org/10.17504/protocols.io.3fxgjpn](https://doi.org/10.17504/protocols.io.3fxgjpn)

Tania Kurbessoian^{1,2}

¹Renata Haro; ²Jason E. Stajich



Tania Kurbessoian



Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.3fxgjpn>

Protocol Citation: Tania Kurbessoian 2019. Cultivating Melanized Fungi from Biological Soil Crust and Rock Surfaces. [protocols.io https://dx.doi.org/10.17504/protocols.io.3fxgjpn](https://dx.doi.org/10.17504/protocols.io.3fxgjpn)

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

Successful Protocol!

Created: May 28, 2019

Last Modified: July 10, 2019

Protocol Integer ID: 23767

Keywords: Melanized fungi, black yeasts, culturing, biological soil crust, rock surfaces, melanized fungi from biological soil crust, melanized fungi, isolating fungi, cultivating melanized fungi, growing fungi, fungi in question, fungi, fungi from the soil, biocrust soils onto microbiological media, strains of organism, microorganism, other microorganism, different microbiological technique, living strains of organism, utilizing different microbiological technique, starting spore, present in the biocrust, biocrust, living strain, microbiological media, plating biocrust soil, organism, life strain, strain, oligotrophic media, soil slurry, soil, antibiotic, culture independent analysis, pathological disease

Abstract

As the interest to understand melanized fungi becomes more of a focus due to pathological diseases, there needs to be a clearer method to isolate and identify the fungi from their surroundings. Culture independent analysis has helped bloom scientist understandings of the vast quantities of microorganisms around us, but culturing and identifying has always been a struggle. We must look into utilizing different microbiological techniques to help better understand, isolate, and to apply Koch's postulates to prove they are the cause of said disease. Growing and isolating fungi has always been an issue especially if the fungi in question is a slow growing fungi, where fast growing fungi or other microorganisms can grow and surpass the field of view and compete with the slow growing fungi. A series of dilutions, antibiotics and oligotrophic media can all counteract the issues to provide you with a clear window to help isolate your fungi in question.

There is still a need to verify that living strains of organisms are actually present in the environment. One approach is to use "culture dependent" method to obtain strains of organisms present in the biocrust. This is achieved by plating biocrust soils onto microbiological media in order to further isolate life strains. Serial dilution is used to isolate the fungi from the soil. This method involves making a soil slurry by diluting the soil sample with increasing higher proportion of water or media in order to get a low starting concentration of starting spores or material growing on the Petri dishes.

Materials

MATERIALS

⊗ Asparagine

⊗ Glucose **P212121 Catalog #Glucose**

⊗ Potassium phosphate (dibasic) **P212121**

⊗ Petri dish, 100×15mm **Fisher Scientific Catalog #FB0875712**

⊗ Falcon® Serological Pipettes, 1 mL 1000 Pipettes **STEMCELL Technologies Inc. Catalog #38001**

⊗ 1.5mL Microtubes


Troubleshooting

- 1 Prepare Glucose Asparagine Agar (GAA)/1L
 0.500g Dibasic Potassium Phosphate
 0.500g Asparagine
 10.00g Glucose
 15.00g Agar

Autoclave

Prepare filtered antibiotics for 1 liter agar solution. Either Gentamycin, Tetracycline, Kanamycin, Streptomycin or Chloramphenicol will work.

2




Visual protocol for clarity.

3

1m



 0.1 g of sample (either biocrust, soil or rock surface can be used) placed in microcentrifuge tube.

3.1 Add  1000 μ L Sterile H₂O into the same tube.

30s

3.2 Label this 10⁰.

30s

4 Vortex (on highest setting) sample for 10 minutes until solution becomes a slurry.


10m

Note

Rock surface samples may not become a slurry.


5 **Preparing the Serial Dilution**

2m

Prepare 3 other microcentrifuge tubes with  900 μ L sterile H₂O .

5.1 Label them 10⁻¹, 10⁻², 10⁻³.

45s

5.2 Take  100 μ L slurry solution from 10⁰ and add it to 10⁻¹. Vortex 10⁻¹.

30s

5.3 Then take  100 μ L 10⁻¹ and add it to 10⁻². Vortex 10⁻².

30s


5.4 Then take  100 μ L 10⁻² and add it to 10⁻³. Vortex 10⁻³.

30s

6 Set up 12 empty Petri Plates and label them with appropriate labels. I.e. Date, Media (list antibiotic used), Crust type, Dilution, Iteration (Plate 1, Plate 2 or Plate 3).

**Note**

For more statistical power we must set them up in triplicate.

6.1 Pipette out  100 μ L 10^0 into first three plates that were properly labeled for 10^0 .

6.2 Pipette out  100 μ L 10^{-1} into next three plates labeled for 10^{-1} .

6.3 Pipette out  100 μ L 10^{-2} into next three plates labeled for 10^{-2} .

6.4 Pipette out  100 μ L 10^{-3} into next three plates labeled for 10^{-3} .

7 Make sure to have media cooled (to  50 °C *) and ready to go at this point.

Note

*which means you can touch it with your hands without burning yourself.

7.1

Add the media slowly to avoid bubbles. Fill until agar closes all gaps on plate. Fill up to three plates at a time.


7.2

When three plates have been filled start to slowly mix the plate and slurry solution. Do a very careful "Figure 8" motion. Make sure to mix slowly and evenly without touching the lid of the plate.


Note

If it does touch, just replace with a new lid from an unused empty plate.

- 7.3 Continue repeating 5.1 and 5.2 until all plates have been filled and are cooling.
- 8 Let plates air dry for 24 hours, then parafilm to allow for long-term storage.
- 9 New fungi should start popping up on undiluted and next dilution plates in the next couple of days. Most dilute plates will also have sparser filamentous fungi growing.
- 9.1 Melanized fungi will take about 1-2 weeks to show up on plates. Make sure to check on the underside of plates for black dots. Those are the fungi you are looking for!



This is an example of the least diluted plate fungal growth.



This is the back of a plate. Notice the little black dots. These are the melanized fungi/yeasts to collect.