



Mar 01, 2022

Version 1

Data normalisation of RT-qPCR data for detection of SARS-CoV-2 in wastewater V.1

DOI

<https://dx.doi.org/10.17504/protocols.io.b4eqqtdw>

Adrian Roberts¹, Zhou Fang¹, Claus-Dieter Mayer¹, Anastasia Frantsuzova¹, Graeme J Cameron², Livia C T Scorza³

¹Biomathematics and Statistics Scotland (BioSS), James Clerk Maxwell Building, Edinburgh, UK.;

²Scottish Environment Protection Agency (SEPA), Strathallan House, Stirling, UK;

³SynthSys and School of Biological Sciences, University of Edinburgh, Edinburgh, UK.

Adrian Roberts: Developed the protocol;

Zhou Fang: Developed the protocol;

Claus-Dieter Mayer: Developed the protocol

Anastasia Frantsuzova: Developed the protocol

Graeme J Cameron: Developed and implemented the protocol

Livia C T Scorza: Curated the protocol



BioRDM Biological Research Data Management

The University of Edinburgh

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS





DOI: <https://dx.doi.org/10.17504/protocols.io.b4eqqtdw>

External link: <https://informatics.sepa.org.uk/RNAmonitoring/>

Protocol Citation: Adrian Roberts, Zhou Fang, Claus-Dieter Mayer, Anastasia Frantsuzova, Graeme J Cameron, Livia C T Scorza 2022. Data normalisation of RT-qPCR data for detection of SARS-CoV-2 in wastewater . **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.b4eqqtdw> Version created by **BioRDM Biological Research Data Management**

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: January 27, 2022

Last Modified: March 01, 2022

Protocol Integer ID: 57520

Keywords: Wastewater-based epidemiology, Infectious diseases, Public health, Covid-19, SARS-CoV-2, RNA viruses, sewage surveillance, normalisation, rna extraction from wastewater, data normalisation of rt, flow for the normalization process, qpcr data for detection, concentration of viral rna, rna value, rna extraction, flow data, flow historical average, data normalisation, normalisation process, qpcr data, using ammonia concentration, daily value of rna copy, raw measurements as gene copy, wastewater, ammonia concentration, population of the catchment area, normalization process, qpcr for detection, parameters such as the flow volume, data as rna copy, predicted ammonia, such as viral rna isolation, catchment area, normalised output, flow volume, amount of flow, viral rna isolation, rna, viral rna, normalization, rna copy, using flow, fluid flow, high volumes of fluid flow, qpcr, detection of sar

Funders Acknowledgements:

CREW (Scotland's Centre of Expertise for Waters)

Grant ID: CD2019_06 Tracking SARS-CoV-2 via municipal wastewater



Abstract

After obtaining the raw measurements as gene copies per litre using RT-qPCR, a normalisation process is required prior to reporting the data as RNA copies per person.

This is because the concentration of viral RNA in wastewater is affected by both the population of the catchment area at each waterworks, as well as the amount of flow into the works. For example, an area with heavy rainfall will have high volumes of fluid flow, which will dilute RNA values.

Therefore, parameters such as the flow volume of wastewater and population size are used to overcome this bias.

Unfortunately, the flow data are not always accessible for all sites. Three methods were developed by Biomathematics and Statistics Scotland (BioSS) to estimate the flow for the normalization process, depending on data availability:

1. Normalisation using ammonia concentration to estimate flow;
2. Normalisation using flow historical average (when data for method 1 are not available);
3. Normalisation using predicted ammonia to estimate flow (when data for methods 1 and 2 are not available).

For all methods, the normalised outputs were reported as a daily value of RNA copies per person.

These methods are described separately in this protocol.

For other methodologies involving SARS-CoV-2 quantification in wastewater, such as viral RNA isolation and RT-qPCR, please refer to:

dx.doi.org/10.17504/protocols.io.bzv5p686 (RNA extraction from wastewater for detection of SARS-CoV-2)


dx.doi.org/10.17504/protocols.io.bzwap7ae (RT-qPCR for detection of SARS-CoV-2 in wastewater)

Decision tree

- 1 Normalisation requires population size and water flow at the collection site.

Depending on the level of information available for particular site and date (such as ammonia content in wastewater or historical flow) different processing paths are available, as described below in the step by step algorithm.

The decision process is also described in the following figure



- 2 If flow data exist for a specific site and date (Flow[site, date]) GO TO normalisation step **8**, otherwise continue.
- 3 If flow data is not available, but measurements of ammonia concentration exist for a specific site and date (Am[site, date]) GO TO prediction step **7**, otherwise continue.
- 4 If there are no measurements for flow or ammonia concentration for a site on any dates GO TO step **5**, or, alternatively, go to step **6** if historical flow is available.

Using historical data to estimate ammonia

5

The missing ammonia values can be inferred from a generalized additive (GAM) spline model which fits seasonal ammonia data.

```
minAm = min(Am)
am_model = gam(log(Am + 1) ~ s(dates, k = 90) + sites)
Am[site, date] = exp(predict(am_model, date, site)) - 1
Am = max(Am, minAm)
```

Observations:

- the smoothing parameter $k=90$ was chosen by preliminary analysis
- the predicted value is prevented from being lower than ever observed

GO TO to flow prediction step **7**

Using only population data to estimate flow

6 If a site has no flow/ammonia data at all, use a national linear model based on population to predict flow

```
flow_model = lm(log10(Flow) ~ logPopulations)
pred = predict(flow_model, site_population)
Flow[site, date] = 10^pred
```

Observations:

- One has to take into account that by using flow historical average, the seasonal aspects are not considered, such as heavier rain fall in particular sites and seasons.

GO TO to normalisation step **8**

Using ammonia data to estimate flow

- 7 When flow measurements are unavailable, but ammonia concentration in wastewater is available (directly or by estimation), a linear mixed statistical model can be used to estimate flow.

More specifically, a random coefficients model is used combining data from ammonia content and the population of each specific site. In this model the log of the flow is regressed on the log of the population and log of ammonia, which are used as fixed effects.

This model is used to calculate site specific coefficients, for those sites which have at least 25 dates with both ammonia and flow data available. For the remaining sites the "fixed" effects are used.

```
flow_model = lmer(logflow ~ logpop + logammonia + (logammonia |  
site), REML=TRUE, data)  
ammoniacoeff[site] = coef(flow_model)[data_rich_sites]  
ammoniacoeff["Unknown"] = fixef(flow_model)
```

The precomputed parameters are used to predict the flow (a table with site specific coefficients is attached to this protocol).

```
param = ammoniacoeff[site] || ammoniacoeff["Unknown"]  
Flow[site, date ] = 10^(param$intercept+param$slope.lgp*  
log10(site_population ) +  
  
param$slope.lgammonia*log10(Am[site,date]))
```

Observations:

- The coefficients parameters were update only every so often, with a degree of care like identifying outliers in the data.

 flow_mixed_model_coefficients_20...

GO TO to normalisation step **8**

Normalisation using flow and population data

- 8 To produce a daily value of RNA copies per person, the raw RNA measurement is multiplied by the daily flow total, and divided by the population served at each site. The flow for a specific site (waterworks location) is either measured directly or estimated using one of the methods from above.

```
normalized = gen_copies*Flow[site, date]/(site_population)/1e6 #  
unit [Mgc/p/d]
```

Code example

- 9 For illustration purposes we present the extract of the code for prediction of ammonia and flow which is used in the BioSS processing pipeline in R.

Command

normalisation - ammonia prediction

```
library(mgcv)

Rdate = as.Date(strptime(RNA$Date, "%m/%d/%Y"))
Rfdate = as.numeric(Rdate)[!is.na(RNA$Ammonia..mg.l.)]
Rfsite = factor(RNA$Site)[!is.na(RNA$Ammonia..mg.l.)]
RfAm = RNA$Ammonia..mg.l.[!is.na(RNA$Ammonia..mg.l.)]

flowobj = gam(log(RfAm + 1) ~ s(Rfdate, k = 90) + Rfsite) #k chosen
by preliminary analysis

# flow ammonia model
wwdat$Flow-> FlowMean
Am = wwdat$Ammonia
Pop = rep((sitedata$PercentageNationalPop * sitedata$NationalPop)
[isite], length(FlowMean))

# 0 = has flow, 1 = has only ammonia, 2 = has neither
dataType = is.na(FlowMean)*(1+ is.na(Am))
# do we have *any* flow/ammonia data for site?
if ((sum(!is.na(wwdat$Flow)) +sum(!is.na(wwdat$Am)) )>0){
  minAm = min(Am, na.rm=T)
  #fill in missing ammonias with ammonia from model
  if (sum(!is.na(Am)) > 0) Am[is.na(Am)] = exp(predict(flowobj,
data.frame(Rfdate = as.numeric(as.Date(strptime(wwdat$Date,
"%m/%d/%Y"))), Rfsite = wwdat$Site))[is.na(Am)]) - 1
  Am = pmax(Am, minAm) #do not allow ammonia to go below
minimum seen

  if (sitename %in% ammoniacoeff$site){
    param = ammoniacoeff[ammoniacoeff$site == sitename,]
    # note adrian is based on log10 values, we reverse
the reparameterisation
    pred =
matrix(rep(10^(param$intercept+param$slope.lgpop*log10(
param$Population )+param$slope.lgammonia*log10
(Am[is.na(wwdat$Flow)])), 3), ncol = 3)
  } else {
    # not in ammonia est, use unknown
    param = ammoniacoeff[ammoniacoeff$site == "Unknown".1
```



```
      pred =  
matrix(rep(10^(param$intercept+param$slope.lgpop*log10( Pop[1]  
)+param$slope.lgammonia*log10 (Am[is.na(wwdat$Flow)])), 3), ncol = 3)  
}  
FlowMean[is.na(wwdat$Flow)] = pred[,1]  
# if prediction fails use mean  
FlowMean[is.na(FlowMean)] = mean(FlowMean, na.rm = T)  
  
} else {  
  # if a site has no flow/am data at all, use a national linear  
  model based on population  
  logP = log10((sitedata$PercentageNationalPop *  
sitedata$NationalPop)[match(RNA$Site, sitedata$Simple)])  
  
  pred = predict(lm(log10(RNA$Flow) ~logP), newdata = list(logP  
= log10(Pop)),interval = "prediction", level = WaterPredLev)  
  FlowMean[is.na(FlowMean)] = 10^pred[,1]  
}  
  
var2 =  
wwdat$N1.Reported*FlowMean/(sitedata$PercentageNationalPop[isite]*site  
data$NationalPop[1]) /1e6 # calc Mgc/p/d
```

