AUG 05, 2020

De-salting of tryptic digested peptides

avinash.kale¹

¹domnic colvin

UM-DAE Centre for Excellence in Basic Sciences

avinash.kale

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjdqki5w

Document Citation: avinash.kale 2020. De-salting of tryptic digested peptides.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bjdqki5w>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Aug 05, 2020

Last Modified: Aug 05, 2020

DOCUMENT integer ID:
 40080

De-salting of tryptic digested peptides

The desalting of tryptic digested samples were optimized and performed using octadecyl carbon chain (C18) pipette tips (ZipTip® Pipette Tips, Millipore). Briefly, for the C18 tips, the desalting pipette tip was pre-equilibrated using a solution of 50% Acetonitrile (SRL) in LC-MS grade water and then equilibrated twice in 0.1% formic acid in LC-MS water. After carefully and gently pipetting the entire sample up and down for at least 10 times, the membrane was washed with 0.1% formic acid in LC-MS water. The peptides were eluted in 70% acetonitrile in LC-MS grade water and vacuum dried and stored for further analysis(26).