



Version 1

Dechoriation of zebrafish embryos with Pronase for metronidazole-mediated β -cell ablation V.1

Sandra Rieger¹

¹University of Miami

dx.doi.org/10.17504/protocols.io.23zgpp6



Lili Liang

ABSTRACT

Summary:

Zebrafish embryos develop in their eggshell (the chorion) until hatching (~48-72 hours post fertilization). Because the chorion is impermeable to water, it needs to be removed prior to metronidazole treatment and β -cell ablation in Tg(*ins:NTR-mCherry*) transgenic zebrafish embryos. This can be achieved either with forceps or by enzymatic removal using Pronase. The earliest time of removal should be after completion of gastrulation to avoid bursting of the yolk and potential infections.

Diabetic Complication:



Neuropathy

DOI

dx.doi.org/10.17504/protocols.io.23zgpp6

EXTERNAL LINK

<https://www.diacomp.org/shared/document.aspx?id=220&docType=Protocol>

PROTOCOL CITATION

Sandra Rieger . Dechoriation of zebrafish embryos with Pronase for metronidazole-mediated β -cell ablation. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.23zgpp6>

KEYWORDS

Dechoriation of zebrafish embryos, NTR-mCherry, Neuropathy

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

May 21, 2019

LAST MODIFIED

Jul 31, 2019

PROTOCOL INTEGER ID

23385

MATERIALS TEXT

MATERIALS

[Pronase from Streptomyces](#)

[griseus Roche Catalog #10165921001](#)

[Sea salt Instant Ocean \(Pet store\)](#)

[1-phenyl-2-thiourea \(PTU\) Sigma](#)

[Aldrich Catalog #P7629](#)

Reagent/Material	Quantity Required
Pronase from <i>Streptomyces griseus</i>	1 g
Sea salt	0.3 g/ liter
1-phenyl-2-thiourea (PTU)	0.003 %

Reagent Preparation:

Reagent 1: Embryo medium

Preparation: Add 0.03 % Instant Ocean salt (Pet store) into double distilled water.

Reagent 2: Pronase stock solution

Preparation: Make a 10 mg/ml stock solution (10 ml) of Pronase and add 1 ml each into ten 15 ml falcon tubes. These can be frozen at -20°C until needed.

Reagent 3: Embryo medium with 1-phenyl-2-thiourea (PTU) to prevent pigment formation.

Preparation: To embryo medium in a small beaker add a final of 0.003% PTU and dissolve overnight using a magnetic stir bar. Keep PTU solution in the dark.

Note:

[Roche, RRID:SCR_001326](#)

[Sigma-Aldrich, RRID:SCR_008988](#)

Protocol:

- 1 For dechoriation, add 9 ml of embryo medium containing 0.003% PTU to 15 ml Falcon tubes containing 1 ml of the Pronase stock solution to make a 10 ml working solution of 1 mg/ml. Pour the working solution into a small petri dish and place the embryos still in their chorion into the solution. Swirl the embryos until the chorion becomes soft (tear off soft chorion with forceps under a stereomicroscope, typically after 3-5 minutes at 24 hours post fertilization). Immediately transfer the embryos into a petri dish with fresh embryo medium. Wash three times in fresh embryo medium.

Potential Pitfalls:

- 2
 1. The embryos do not survive after treatment: This could be if embryos were incubated too long in Pronase. Decrease the incubation time and increase the number of washes.

2. Dechoriation does not work: Pronase might be too old or animals are older than 24 hours post fertilization.

Increase incubation time in Pronase until chorions become soft. This can be up to 10 minutes in 48 hpf larvae.