VERSION 1

JUL 20, 2018

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.qnndvde](https://doi.org/10.17504/protocols.io.qnndvde)

External link:

<https://doi.org/10.1371/journal.pone.0200523>

Protocol Citation: Stefano Orlando, Isotta triulzi, Fausto Ciccacci, Ilaria Palla, Leonardo Palombi, Maria Cristina Marazzi, Marina Giuliano, Marco Floridia, Sandro Mancinelli, Elsa Mutemba, Giuseppe Turchetti 2018. Delayed diagnosis and treatment of tuberculosis in HIV+ patients in Mozambique: a cost-effectiveness analysis of screening protocols based on four symptom screening, smear microscopy, urine LAM test and Xpert MTB/RIF.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.qnndvde>

Delayed diagnosis and treatment of tuberculosis in HIV+ patients in Mozambique: a cost-effectiveness analysis of screening protocols based on four symptom screening, smear microscopy, urine LAM test and Xpert MTB/RIF V.1

PLOS One

Stefano Orlando¹, Isotta triulzi², Fausto Ciccacci¹, Ilaria Palla², Leonardo Palombi¹, Maria Cristina Marazzi³, Marina Giuliano⁴, Marco Floridia⁴, Sandro Mancinelli¹, Elsa Mutemba⁵, Giuseppe Turchetti²

¹University of Roma "Tor Vergata";

²Institute of Management, Scuola Superiore Sant' Anna;

³LUMSA University, Rome;

⁴National Center for Global Health, Istituto Superiore di Sanità, Rome;

⁵DREAM programme, Comunità di Sant'Egidio

Stefano Orlando

ABSTRACT

BACKGROUND

Tuberculosis (TB) represents the ninth leading cause of death worldwide. In 2016, are estimated 1.3 million TB deaths among HIV negative people and an additional 374,000 deaths among HIV positive people. In 2016, are estimated 1.4 million new cases of TB in people living with HIV (PLWH), 74% of whom were living in Africa. In light of these data, the reduction of mortality caused by TB in PLWH is strongly required specially in low-income countries as Mozambique. According to international guidelines, the initial TB screening in HIV+ patients should be done with the four symptoms screening (4SS: fever, current cough, night sweats and weight loss). The diagnostic test more used in resource-limited countries is smear microscopy (SMEAR). World Health Organization (WHO) recommended Lateral Flow urine LipoArabinoMannan assay (LF-LAM) in immunocompromised patients; in 2010 WHO endorsed the use of Mycobacterium Tuberculosis/Rifampicin (MTB/RIF) test for rapid TB diagnosis but the assay is not used as screening test in all HIV+ patients irrespectively of symptoms due to cost and logistical barriers.

The paper aims to evaluate the cost-effectiveness of three screening protocols: standard (4SS and SMEAR in positive patients to 4SS); *MTB/RIF*, *LF-LAM* / *MTB/RIF*.

METHODS

We developed a model to assess the cost-effectiveness of the *MTB/RIF* protocol versus the common *standard* and *LF-LAM* / *MTB/RIF* protocol. The model

MANUSCRIPT CITATION:

Orlando S, Triulzi I, Ciccacci F, Palla I, Palombi L, Marazzi MC, Giuliano M, Floridia M, Mancinelli S, Mutemba E, Turchetti G (2018) Delayed diagnosis and treatment of tuberculosis in HIV+ patients in Mozambique: A cost-effectiveness analysis of screening protocols based on four symptom screening, smear microscopy, urine LAM test and Xpert MTB/RIF. PLoS ONE 13(7): e0200523. doi: [10.1371/journal.pone.0200523](https://doi.org/10.1371/journal.pone.0200523)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jun 01, 2018

Last Modified: Jul 20, 2018

PROTOCOL integer ID:
12718

Keywords: cost-effectiveness; tuberculosis; Xpert MTB/RIF; four symptoms screening, smear microscopy, LF-LAM, HIV/AIDS, delayed diagnosis and treatment, new TB infections, low income country

considered a sample of 1,000 HIV+ antiretroviral treatment naïve patients in Mozambique. We evaluated disability-adjusted life year (DALY) averted for each protocol, cost per DALY, and incremental cost-effectiveness ratio (ICER), over 1-year, assuming a national healthcare system perspective. The model considered the delayed diagnosis and treatment as the time elapsed between the a false negative test and the diagnosis of TB. On top of that there is a possible health system delay defined as the time interval between positive test and treatment initiation due to the a delay in the delivery of results due to weaknesses in the organization of services. We conducted a sensitivity analysis on more relevant variables.

RESULTS

The MTB/RIF protocol was cost-effective as compared to the standard protocol with an ICER of \$56.54 per DALY saved. In a cohort of 1,000 patients MTB/RIF and LF-LAM / MTB/RIF protocol generated 1,281 and 1,254 DALY's saved respectively, with a difference of 174 and 147 DALY respect to the standard protocol.

The total cost of MTB/RIF protocol was lower (\$92,263) than the standard (\$147,226) and the LF-LAM / MTB/RIF protocol (\$113,196).

Therefore, the cost per DALY saved including new infections due to delayed diagnosis and treatment with the standard protocol was \$79.06, about 5 fold higher than MTB/RIF and LF-LAM / MTB/RIF protocols.

The cost of new and additional TB infections due to delays in diagnosis plus health system delay seemed the more relevant costs in our analysis. The low sensibility and sensitivity of the standard protocol led to a high number of false negatives, thus delayed TB diagnoses and treatment lead to the development of newly transmitted TB infections.

CONCLUSIONS

Our study shows that the MTB/RIF adoption could lead an increasing of TB case-finding and a reduction in costs compared with standard and LF-LAM / MTB/RIF protocols

ATTACHMENTS

[Data_rev2.xlsx](#)