SEP 25, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bj8rkrv6

Protocol Citation: Diep R Ganguly, Anna Behle 2020. Denaturing PAGE for resolving RNA. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bj8rkrv6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 24, 2020

Last Modified: Sep 25, 2020

PROTOCOL integer ID:
 40945

Keywords: RNA, Electrophoresis, Denaturing-PAGE, PAGE

Denaturing PAGE for resolving RNA

Forked from [Denaturing RNA Urea-PAGE](#)

Diep R Ganguly¹, Anna Behle²

¹University of Pennsylvania; ²Institute for Synthetic Microbiology

Pogson Group

Diep R Ganguly

University of Pennsylvania, The Australian National Universi...

ABSTRACT

Protocol denaturing PAGE separation and identification of total RNA extracts using Urea-PAGE with the BioRad Mini-PROTEAN II / Tetra Cell electrophoresis system. Modified from dx.doi.org/10.17504/protocols.io.gszbwf6.

GUIDELINES

Use filter tips and ensure you are using RNase free components and surfaces.

This is a fantastic resource to help planning and troubleshooting: RNA: A Laboratory Manual (https://www.cshlpress.com/default.tpl?cart=1598856781901971730&fromlink=T&linkaction=full&linksortby=oop_title&eqSKUdataarq=725).

Also see [10.3791/1485](https://doi.org/10.3791/1485).

MATERIALS

MATERIALS

- ✂ ssRNA Ladder - 25 gel lanes **New England Biolabs Catalog #N0362S**
- ✂ dsRNA Ladder - 25 gel lanes **New England Biolabs Catalog #N0363S**
- ✂ Low Range ssRNA Ladder - 25 gel lanes **New England Biolabs Catalog #N0364S**
- ✂ SYBR Gold **Thermo Scientific Catalog #S-11494**
- ✂ TEMED **Bio-rad Laboratories Catalog #1610801**
- ✂ Mini Protean tetra cell **Bio-rad Laboratories**
- ✂ SYBR Green **Thermo Fisher Scientific**
- ✂ Ethidium bromide [EB, EtBr] **Bio Basic Inc. Catalog #EB0195.SIZE.1g**
- ✂ DEPC Water **Bio Basic Inc. Catalog #DD1005(D0121).SIZE.4x500ML**
- ✂ Axygen 0.5 mL PCR tubes, 0.5 mL, thin wall, clear, flat caps **Corning Catalog #PCR-05-C**
- ✂ Thermal cycler **Contributed by users**
- ✂ RNaseZap® RNase Decontamination Solution **Life Technologies Catalog #AM9780**
- ✂ Filter tips **Contributed by users**
- ✂ Depc (Diethyl Pyrocarbonate) **VWR international Ltd Catalog #80730-888**
- ✂ 40% Acrylamide-bisacrylamide (19:1) **Contributed by users**
- ✂ 30% Acrylamide / Bis. solution **Bio-rad Laboratories Catalog #161-0158**
- ✂ 0.1% SDS solution **Contributed by users**
- ✂ 10% Ammonium persulfate (APS) **Sigma Aldrich**
- ✂ Gel Loading Buffer II (Denaturing PAGE) **Thermo Fisher Catalog #AM8546G**
- ✂ TBE (10X), RNase-free **Thermo Fisher Catalog #AM9865**
- ✂ Mini-PROTEAN® Spacer Plates with 0.75 mm Integrated Spacers **BioRad Sciences Catalog #1653310**
- ✂ Mini-PROTEAN® Short Plates **BioRad Sciences Catalog #1653308**

SAFETY WARNINGS

- ! Wear vinyl or nitrile gloves when handling PAA (neurotoxic).
Make sure to thoroughly wash wells before loading samples.
2x RNA loading buffer contains formamide.

BEFORE START INSTRUCTIONS

Thoroughly clean all components with 70% ethanol and RNase away (or 0.05 - 0.1 % SDS). Ensure RNA samples of interest have been normalized to a known concentration with nuclease-free/DEPC-treated water.

Gel preparation

- 1 Before preparing the gel, clean all components, gel tank, and surfances with 70 % ethanol (can use 0.1% SDS or RNase Zap to be extra thorough, we do this periodically).
- 2 Assemble gel plates in the green brackets and place on casting unit. Ensure glass plates are not cracked and are sitting flush against the separator plate and on the rubber mats (to prevent gel leakage).
- 3 Use the following table to prepare the required gel recipe using either 30 % *or* 40 % PAA (~4 mL=1x0.75 mm gel).

PAA percentage	8 %	10 %	12 %	15 %
30 % / 40 % PAA	2 / 2.67 mL	2.5 / 3.33 mL	3 / 4 mL	3.75 / 5 mL
8 M Urea	5 g	5 g	5 g	5 g
10x TBE	1 mL	1 mL	1 mL	1 mL
10 % APS	80 µL	80 µL	80 µL	80 µL
TEMED	10 µL	10 µL	10 µL	10 µL
DEPC-treated H ₂ O	ad 10 mL	ad 10 mL	ad 10 mL	ad 10 mL

Urea-PAGE recipe

- 4 Dissolve urea in 10x TBE, PAA, and 1/2 volume DEPC-treated H₂O.

Warming solution (e.g. 37°C water bath) with gentle agitation will help to dissolve urea.

Safety information

Wear vinyl or nitrile gloves and work in fumehood when handling PAA (neurotoxic!)

- 5 Once dissolved, allow to cool to room temperature and add DEPC-treated H₂O up to 10 mL.

- 6 Add 80 μL APS (10 %) and 10 μL TEMED to the gel solution and gently swirl once. Immediately pipette solution into gel apparatus (~ 4.5 mL, fill until just before it overflows to prevent bubbles when inserting comb). Retain some gel mixture to check for complete polymerisation.
- 7 Immediately insert comb tilted at an angle (to minimise trapping air bubbles). Make sure you are wearing appropriate gloves. Lower completely and allow gel to polymerise for >15 mins. Gels can be stored at 4°C wrapped in plastic (use with 3-5 days).

RNA preparation

- 8 Determine desired loading amount of RNA per well (e.g. 0.5 - 2 μg RNA). Normalize RNA concentrations accordingly (e.g. 100 - 200 ng/ μL).
- 9 Add required RNA to equal volume 2x gel loading buffer II (Ambion). Up to to 50% can be RNA sample although less is better. Final volumes for loading may vary, however, less than 5 μL results in poor banding when using the Mini protean system (10 μL / gel works well). Keep prepared samples @ 4°C until ready for denaturation and loading.

Safety information

RNA loading dye contains formamide. Wear goggles/lab coat/ gloves!

RNA loading

- 10 Once remaining gel solution has polymerised, remove from casting bracket and insert into running bracket (keep combs). Fill interior with 1x TBE and ensure there are no leaks (~ 5 mins). If leaking, re-adjust and check again. Place bracket into gel tank and fill with 1x TBE.
- 11 Remove combs and rinse individual wells thoroughly.

- 12 Connect electrodes, and pre-run gel @ 10 mA / gel for ~20-30 minutes to remove residual urea.
- 13 During the pre-run, denature RNA samples and ladder, in gel loading buffer II, at 72°C for 5 min. Once finished, immediately keep at 4°C until ready to load (minimise time kept at 4°C).
- 14 Turn power supply off and rinse the well a second time.
- 15 Pipette denatured samples into wells slowly while ensuring there are no air bubbles trapped in the pipette tip (Ambion recommends loading while samples are still hot).
- 16 Close the lid and plug electrodes into power supply. Separate RNA @ 10 mA / gel for ~1-2 hours (make sure you have a reliable power supply that holds current, particularly as urea has poor conductivity). Allowing the upper marker, from the Ambion loading buffer II, to run off the gel is a reasonable indicator for running time.

Visualization

- 17 After sufficient running time switch off power supply, pour out buffer, disassemble gel and clean components. Transfer gel to clean container and rinse once with deionized water.
- 18 Incubate gel in 0.5 µg ethidium bromide / mL 1x TBE for ~15 minutes (or use SYBR gold/green safe stains). Rinse 3x with water and allow to destain for ~2 minutes. Visualize.