APR 03, 2019

DENV Titration

Fabio Gomes¹

¹Johns Hopkins University

Fabio Gomes

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.zsdf6a6

Protocol Citation: Fabio Gomes 2019. DENV Titration. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.zsdf6a6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Apr 03, 2019

Last Modified: Apr 03, 2019

PROTOCOL integer ID:
22053

Fixation of Cells

- 1 Prepare fixative - 80% methanol in water

2 Dump Methylcellulose overlay and blot on paper towels

3 Wash gently with 1X PBS. Incubate for 10 min. Blot and dry. ⌚ 00:10:00

4 Add 0.5 mL fixative to each well. Allow plates to sit at room temperature for 10 min.

Note

Note: Fixed cells can be stored at -70oC for future use. Leave methanol on cells when freezing.

Antibody

5 Wash plates with PBS. Incubate for 10 min. ⌚ 00:10:00

Note

If frozen plates are being used, incubate at 37oC for 30 min.

6 Prepare antibody dilutions in 5% skim milk, PBS [M] 5 Mass / % volume Skim Milk

Note

4G2 1:2000

7 Add 200 µl to each well. Incubate at 37°C for 1 hour on a rocker. 01:00:00

37 °C

Minimize contact with each well by maintaining contact of the pipette tip with the wall of well.

8 Dump off primary antibody solution and tap plates on paper towels to remove excess solution

9 Wash with 1 mL 5% skim milk, PBS

10 Prepare antibody dilutions in 5% skim milk, PBS.

Note

Anti-mouse-Per 1:2,000

11 Add 200 µl to each well. Incubate at 37°C for 1 hour on a rocker. 01:00:00 37 °C

Color Development

12 Dump off antibody solution. Tap on paper towels

13 Wash twice with PBS. Tap on paper towel

14 Add 160 µl of TrueBlue substrate per well

15 Place on rocker at RT until plaques develop (10 min or longer).

16 Dump off peroxidase substrate.

17 Count plates or incubate at 4°C for up to 7 days.