AUG 15, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjspkndn

Protocol Citation: Angel A Justiz-Vaillant 2020. Detection of anti- keyhole limpet hemocynin (anti-KLH) antibodies by double immunodiffusion (Ouchterlony) technique.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjspkndn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 15, 2020

Last Modified: Aug 15, 2020

PROTOCOL integer ID:
 40495

Detection of anti- keyhole limpet hemocynin (anti-KLH) antibodies by double immunodiffusion (Ouchterlony) technique.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

ABSTRACT

Keyhole limpet hemocyanin (KLH) is a cooper-containing protein comprising of subunits with MW of 400 kDa. This protein is found in the hemolymph of the sea mollusk *Megathura crenulata*. It has the ability to enhance the host's immune response by interacting with monocytes, T cells and macrophages. KLH has been used primarily as a carrier for vaccines and antigens [1]. It was found that chicken immunized with KLH bound peptide raised an anti-KLH immunoresponse [2]. This can be tested by a single method such as the Ouchterlony technique.

Reference

1. Aarntzen EH, de Vries IJ, Göertz JH, et al. Humoral anti-KLH responses in cancer patients treated with dendritic cell-based immunotherapy are dictated by different vaccination parameters. *Cancer Immunol Immunother*. 2012;61(11):2003-2011. doi:10.1007/s00262-012-1263-z
2. Justiz Vaillant AA, Anderson MF, Smikle M, Wisdom B, Mohammed W, et al. (2013) Development of Anti HIV Gp120 and HIV Gp41 Peptide Vaccines. *J Vaccines Vaccin* 4: 206. doi: 10.4172/2157-7560.1000206

MATERIALS

MATERIALS

10mg KLH (Keyhole Limpet Hemocyanin) (Immunological Grade) G-
Biosciences Catalog #786-088

- 1 Detection of anti-keyhole limpet hemocynin antibodies by double immunodiffusion is carried out.

- 2 Briefly, 1% agarose gels are prepared and wells cut into the gel using a template.
- 3 Initially, aliquots of 25 µl each of KLH in concentration of 1 mg/ml are applied to the centre well.
- 4 The peripheral wells are filled with 25 µl of isolated chicken IgY (1 mg/ml) in PBS pH 7.4.
- 5 The gels are incubated at RT for 48–72 hours.
- 6 After that the gels are examined for precipitin lines.
- 7 An anti-KLH developed in cat is included as positive control and turtle serum as a negative control.
- 8 The positive results are taken as the presence of precipitin line/s and negative results, the absence of precipitin lines.