VERSION 2

MAR 28, 2019

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.wyqffvw](https://doi.org/10.17504/protocols.io.wyqffvw)

External link:

<https://doi.org/10.1371/journal.pone.0213493>

Protocol Citation: Cecilia Monica Rodriguez García, Leticia LPE Peraza Echeverría 2019. Detection of anti-phytopathogenic fungal activity. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.wyqffvw>

MANUSCRIPT CITATION:

Rodríguez-García CM, Ruiz-Ruiz JC, Peraza-Echeverría L, Peraza-Sánchez SR, Torres-Tapia LW, Pérez-Brito D, Tapia-Tussell R, Herrera-Chalé FG, Segura-Campos MR, Quijano-Ramayo A, Ramón-Sierra JM, Ortiz-Vázquez E (2019) Antioxidant, antihypertensive, anti-hyperglycemic, and antimicrobial activity of aqueous extracts from twelve native plants of the Yucatan coast. PLoS ONE 14(3): e0213493. doi: [10.1371/journal.pone.0213493](https://doi.org/10.1371/journal.pone.0213493)

Detection of anti-phytopathogenic fungal activity V.2

PLOS One

Cecilia Monica Rodriguez García¹, Leticia LPE Peraza Echeverría¹

¹Centro de Investigación Científica de Yucatán

Leticia LPE Peraza Echeverría

Centro de Investigación Científica de Yucatán

ABSTRACT

Here we describe the detection of in vitro anti-phytopathogenic fungal activity using the broth dilution method to find the MIC (minimum inhibitory concentration) of an aqueous extract of leaves. Four concentrations were assayed: 1, 2.5, 5, and 10% (v/v) in PDB medium. Each concentration was inoculated with a spore suspension of the selected fungus. PDB without spores was used as negative control and with the addition of a commercial fungicide like positive control. After the inoculation of the fungus, the tubes were incubated for three days in the dark at 26-28 °C. After the incubation time, the tubes were checked for growth of the fungus, if no growth is observed, 25 µL of the suspension are transferred to PDA medium in Petri dishes of 3 cm diameter. Evaluations were made on day 7. The MIC was determined as the lowest concentration of aqueous extract preventing the growth of macroscopically visible mycelium.

GUIDELINES

- 35 x 10 mm polystyrene Petri dish from Corning
- Commercial fungicide (Miconazole nitrate, 20 mg/L)
- Culture room for the fungal growth (26 ± 2 °C at 12/12 h photoperiod)
- Microscope Axioplan from Zeiss
- Everything must be done in a laminar flow hood

MATERIALS

MATERIALS

☒ Potato Dextrose Agar, Difco **Fisher Scientific Catalog #DF0013-15-8**

☒ Potato Dextrose Broth, Difco **Fisher Scientific Catalog #DF0549-17-9**

BEFORE START INSTRUCTIONS

- Petri dishes with PDA medium
- PDB medium
- Everything should be sterilized

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jan 10, 2019

Last Modified: Mar 28, 2019

PROTOCOL integer ID:
19184

Keywords: bioassay, in vitro detection, aqueous plant extracts

Dilution of the aqueous extract of leaves

- 1 ■ Performs three dilutions from the 10 % aqueous extract (AE)

Dilution number	AE (μL)	CS (μL)	H ₂ O (μL)	final volume (μL)	concentration (%)
1	500	100	400	1000	5
2	250	100	650	1000	2.5
3	125	100	775	1000	1.25
	---	---	---	1000	10

CS- conidial suspension (200 conidium/μL)

Note

Do not add de CS, only make the mix of AE and water, CS is added tube by tube

Dilution of the aqueous extract of leaves

2

1. Add to a sterile 1.5 mL eppendorf tube mix 1 and then 12.5 μL of CS (in duplicate)
2. Repeat num 1 for the next concentration and so on..
3. For the 10% AE just put in each tube 125 μL 10% AE plus 12.5 μL of CS, (in duplicate)
4. The positive control is prepared by adding 1 μL of a commercial fungicide to 125 μL of water

- and 12.5 µL of CS (in duplicate)
5. Two negative controls: one is only 125 µL of water and the other is 125 µL of PDB and both with 12.5 µL of CS (in duplicate)

Bioassay conditions

- 3 Once the conidial suspension has been added, homogenize very well, seal the tubes and leave them in the dark at a temperature of 26-28 ° C for 3 days.

- 4 After three days, 25 µL of the suspension is transferred in the center of the PDA medium (polystyrene dish, Corning), in duplicate.

Note

The suspension of each eppendorf tube must be perfectly homogenized

- 5
- Petri dishes are incubated at 26 ± 2 °C at 12/12 photoperiod for 7 days (two replicates /concentration)
 - Evaluation are made on day 7

Expected result

If the aqueous extract of leaves is fungicidal there should be no growth on the PDA medium, if the extract is not fungicidal then there must be growth of the fungus on the PDA medium