



Aug 15, 2020

## Detection of anti- SpA antibodies in egg white tested by double immunodiffusion (Ouchterlony) technique.

DOI

[dx.doi.org/10.17504/protocols.io.bjsqkndw](https://dx.doi.org/10.17504/protocols.io.bjsqkndw)

Angel A Justiz-Vaillant<sup>1</sup>

<sup>1</sup>University of the West Indies St. Augustine

University of the West In...

angel.vaillant@sta.uwi.e...



Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.bjsqkndw](https://dx.doi.org/10.17504/protocols.io.bjsqkndw)

**Protocol Citation:** Angel A Justiz-Vaillant 2020. Detection of anti- SpA antibodies in egg white tested by double immunodiffusion (Ouchterlony) technique.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjsqkndw>

**License:** This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** August 15, 2020

**Last Modified:** August 15, 2020

**Protocol Integer ID:** 40496



## Abstract

Keyhole limpet hemocyanin (KLH) is a copper-containing protein comprising of subunits with MW of 400 kDa. This protein is found in the hemolymph of the sea mollusk *Megathura crenulata*. It has the ability to enhance the host's immune response by interacting with monocytes, T cells and macrophages. KLH has been used primarily as a carrier for vaccines and antigens [1]. It was found that chicken immunized with KLH bound peptide raised an anti-KLH immunoresponse [2]. This can be tested by a single method such as the Ouchterlony technique.

## Reference

1. Aarntzen EH, de Vries IJ, Göertz JH, et al. Humoral anti-KLH responses in cancer patients treated with dendritic cell-based immunotherapy are dictated by different vaccination parameters. *Cancer Immunol Immunother*. 2012;61(11):2003-2011. doi:10.1007/s00262-012-1263-z
2. Justiz Vaillant AA, Anderson MF, Smikle M, Wisdom B, Mohammed W, et al. (2013) Development of Anti HIV Gp120 and HIV Gp41 Peptide Vaccines. *J Vaccines Vaccin* 4: 206. doi: 10.4172/2157-7560.1000206

## Materials

### MATERIALS

 10mg KLH (Keyhole Limpet Hemocyanin) (Immunological Grade) **G-Biosciences Catalog #786-088**



- 1 Detection of anti-keyhole limpet hemocynin antibodies by double immunodiffusion is carried out.
- 2 Briefly, 1% agarose gels are prepared and wells cut into the gel using a template.
- 3 Initially, aliquots of 25  $\mu$ l each of KLH in concentration of 1 mg/ml are applied to the centre well.
- 4 The peripheral wells are filled with 25  $\mu$ l of isolated chicken IgY (1 mg/ml) in PBS pH 7.4.
- 5 The gels are incubated at RT for 48–72 hours.
- 6 After that the gels are examined for precipitin lines.
- 7 An anti-KLH developed in cat is included as positive control and turtle serum as a negative control.
- 8 The positive results are taken as the presence of precipitin line/s and negative results, the absence of precipitin lines.