Jun 25, 2017





Forked from <u>Detection of protein-synthesizing microorganisms in the environment via bioorthogonal non-</u> canonical amino acid tagging (BONCAT)

DOI

dx.doi.org/10.17504/protocols.io.imccc2w

Roland Hatzenpichler





DOI: dx.doi.org/10.17504/protocols.io.imccc2w

External link: <u>http://nebula.wsimg.com/b046306c6f40dd35b12d1819a48c7ce9?</u> AccessKeyId=48B9AC89493A56C67DF6&disposition=0&alloworigin=1

Protocol Citation: Roland Hatzenpichler: Detection of protein-synthesizing microbes via bioorthogonal non-canonical amino acid tagging (BONCAT). protocols.io <u>https://dx.doi.org/10.17504/protocols.io.imccc2w</u>

License: This is an open access protocol distributed under the terms of the <u>Creative Commons Attribution License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: June 25, 2017

Last Modified: March 07, 2018

Protocol Integer ID: 6532

Keywords: bioorthogonal noncanonical amino acid tagging, different bioorthogonal amino acid, detection of protein, synthesizing microorganisms in the environment, noncanonical amino acid, synthesizing microbe, synthesizing microorganism, incorporation of these noncanonical amino acid, new protein, homopropargylglycine, protein, boncat experiments with environmental sample, boncat experiment, lipid microbiology protocol, taxonomic identification of cell, taxonomic identification, whole cells via copper, boncat, azidohomoalanine

Abstract

This protocol provides a detailed description of how to design and perform BONCAT experiments using two different bioorthogonal amino acids, L-azidohomoalanine (AHA) and L-homopropargylglycine (HPG), which are both surrogates of L-methionine. It illustrates how incorporation of these noncanonical amino acids into new proteins can be detected in whole cells via copper-catalyzed or strain-promoted azide-alkyne click chemistry and outlines how the visualization of translational activity can be combined with the taxonomic identification of cells via FISH. Last, the protocol discusses potential problems that might be encountered during BONCAT experiments with environmental samples and how they can be overcome.

Book chapter describing this BONCAT protocol is published in:

Detection of Protein-Synthesizing Microorganisms in the Environment via Bioorthogonal Noncanonical Amino Acid Tagging (BONCAT)

Authors: Roland Hatzenpichler and Victoria J. Orphan

T.J. McGenity et al. (eds.), Hydrocarbon and Lipid Microbiology Protocols, Springer Protocols Handbooks, DOI 10.1007/8623_2015_61, © Springer-Verlag Berlin Heidelberg 2015

Before start

There are two variants of azide-alkyne click reactions:

(1) a Cu (I)-catalyzed cycloaddition reaction and

(2) a strain-promoted version that exploits the high reactivity of a cyclooctyne system that allows the click reaction to take place in the absence of a catalyst.

Both labeling reactions are simple with only a few inexpensive chemicals required. The protocol can be preformed within 1–2 hours to identify transcriptionally active microorganisms in a variety of environmental samples after incubation with a methionine surrogate. In our experience, the BONCAT reaction is generally not influenced by the presence of complex organic (e.g., an extracellular matrix) or inorganic (e.g., minerals or sediment particles) substances. This results in a low level of background noise when applied to environmental samples. While copper-catalyzed click chemistry can be employed to detect both AHA- and HPG-tagged proteins, the strain-promoted click reaction is restricted to the visualization of AHA uptake (because HPG does not contain an azide group).

Solutions required for BONCAT labeling (Unless indicated otherwise, all reagents can be stored at room temperature).

For L-2-amino-4-azidobutanoic acid (L-azidohomoalanine, AHA): Dissolve in nano-pure water, adjust to pH 7.0, filter sterilize (0.2 μm), and store in the dark at 4°C. Prepare stock solutions (typically in the range between 1–100 mM).

For L-2-amino-5-hexynoic acid (L-homopropargylglycine, HPG): Dissolve in nano-pure water, adjust to pH 7.0, filter sterilize (0.2 μ m), and store in the dark at 4°C. Prepare stock solutions (1–100 mM).

Once samples have been incubated with either HPG or AHA, paraformaldehyde fixation (PFA) and preparation of microbial cells on a teflon coated slide or polycarbonate filter should be done following a standard protocol for FISH (e.g. Ishii et al. 2004 or Daims et al 2005). After PFA fixed samples are prepared, the copper-catalyzed (Cu I) click reaction for fluorescence detection of newly synthesized proteins in whole cells requires the following solutions:

Phosphate-buffered saline (PBS): 130 mM NaCl, 5% (v/v) phosphate buffer in nano-pure water, adjust to pH 7.4, and sterilize filter (0.2 μ m).

20 mM Copper sulfate solution (CuSO4 🛿 5 H2O) in nano-pure water, sterilize filter (0.2 μm), and store at room temperature or 4°C.

50 mM THPTA : Tris[(1-hydroxypropyl-1H-1,2,3-triazol-4-yl)methyl]amine in nano-pure water, sterilize filter (0.2 μ m), and store in aliquots at [-20°C.

100 mM freshly prepared sodium ascorbate in 1X PBS (this reducing agent is added to ensure the copper sulfate solution has a sufficient amount of copper I to catalyze the azide-alkyne reaction). Should be made fresh each time and used the same day.

100 mM Aminoguanidine hydrochloride in 1X PBS (this reagent should also be made fresh each time)

1–10 mM stock solutions in either dimethylsulfoxide (DMSO) or dimethylformamide (DMF) for preparation of the dye solution following manufacturer's instructions (Alexa dyes or Cy dyes) for HPG incubations, use azide-conjugated fluorophores for AHA incubations, use alkyne-conjugated fluorophores

See following references for additional information on copper-catalyzed and strain promoted Click and FISH sample preparation.

Ishii, K., M. Mußmann, B. J. MacGregor, and R. Amann. 2004. An improved fluorescence in situ hybridization protocol for the identification of bacteria and archaea in marine sediments. FEMS Microbiology Ecology 50:203-212.

Daims H, Stoecker K, Wagner M (2005) Fluorescence in situ hybridization for the detection of prokaryotes. In: Osborn AM, Smith CJ (eds) Molecular microbial ecology. Bios Advanced Methods, Abingdon, pp 213–239

Bagert JD, Xie YJ, Sweredoski MJ, Qi Y, Hess S, Schuman EM, Tirrell DA (2014) Quantitative, time-resolved proteomic analysis by combining bioorthogonal noncanonical amino acid tagging and pulsed stable isotope labeling by amino acids in cell culture. Mol Cell Proteomics 13(5):1352–1358.

Dieterich DC, Link AJ, Graumann J, Tirrell DA, Schuman EM (2006) Selective identification of newly synthesized proteins in mammalian cells using bioorthogonal noncanonical amino acid tagging (BONCAT). Proc Natl Acad Sci U S A 103:9482–9487

Torne CW, Christensen C, Meldal M (2002) Peptidotriazoles on solid phase: [1,2,3]- friazoles by regiospecific copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides. J Org Chem 67:3057–3064

Agard NJ, Prescher JA, Bertozzi CR (2004) A strain-promoted [3 + 2] azide-alkyne cycloaddition for covalent modification of biomolecules in living systems. J Am Chem Soc 126:15046–15047

Incubation with Bioorthogonal Amino Acid

1 Directly add AHA or HPG using a sterile-filtered (0.2 μM), pH- adjusted (pH 7.0) stock solution yielding a final concentration of 1 nM to 1 mM. Additionally, perform replicate experiments and include replicated incubations without AHA/HPG.

For L-2-amino-4-azidobutanoic acid (L-azidohomoalanine, AHA): Dissolve in nano-pure water, adjust to pH 7.0, filter sterilize (0.2 μ m), and store in the dark at 4 C. Prepare stock solutions of 1–100 mM.

For L-2-amino-5-hexynoic acid (L-homopropargy/glycine, HPG): Dissolve in nano-pure water, adjust to pH 7.0, filter sterilize (0.2 μ m), and store in the dark at 4 C. Prepare stock solutions of 1–100 mM.

Note

The Hatzenpichler lab highly recommends to buy bioorthogonal amino acids from clickchemistrytools. When compared to other companies, like Invitrogen or Sigma, clickchemistrytools products are ~20x cheaper, as similar or better quality.

Cell Fixation

2 Click staining can be performed on living (non-fixed) cells. However, for long term preservation of cells, cell fixaton with formaldehyde, glutaraldehyde, or ethanol is recommended.

To achieve cell preservation for storage at -20C, fix cells according to standard protocols [34] immediately after sampling either by (1) fixation in 3% formaldehyde (PFA) in PBS **(See Steps 3-6)** or (2) by resuspending pelleted biomass in a 1:1 mix of PBS:EtOH. **(See Step 7)**

Note

If working with marine microorganisms, use of 3X PBS is recommended.

Cell Fixation (PFA ONLY)

3 For fixation with PFA, pellet the biomass, remove the supernatant (SN), and resuspend cells in 3% PFA in PBS. For aqueous samples, directly add PFA to reach a final concentration of 3% PFA.

- 4 Fix for either 3 h on ice or 1 h at RT.
- 5 After fixation, Pellet the biomass by centrifugation or filter onto 0.2 µm filters.
- 6 Wash with PBS to remove remaining PFA before resuspending biomass in 1:1 PBS: EtOH. Store at -20°C. Make sure to deposit PFA in the chemical waste.

Cell Fixation (EtOH ONLY)

7 For EtOH fixation, pellet biomass, remove supernatant, resuspend in 1:1 PBS:EtOH, and store at -20°C.

Preparing the Samples for Click Labeling

- 8 Immobilize biomass either on glass slides or filters. Dry at 46°C or, if not available, at 37°C or RT.
- 9 Dehydrate and permeabilize cells by sequentially placing slides or filters for 3 min into 50 mL tubes that contain 50, 80, and 96% ethanol. Dry biomass using pressurized air.
- 10 Pellet sample via centrifugation (16,100g or max. setting for 5 min at RT) and resuspend in 250 μL 80% EtOH.
- 11 Mix by vortex and incubate for 3 min at RT.
- 12 Add 1.5 mL 96% EtOH, mix by vortex, and incubate for 3 min at RT.
- 13 Afterwards, pellet sample via centrifugation and resuspend in 221 µL PBS. Removing small volumes of leftover EtOH is not necessary as it does not interfere with the click reaction.

Cu(I) Click Chemistry

14 If using immobilized biomass, after dehydration of the sample, prepare the dye premix by mixing 1.25μL of 20 mM CuSO₄ solution with 2.50μL of 50 mM THPTA and 0.30μL of alkyne dye. Allow to react for 3 min at RT in the dark. We recommend to perform Cu(I)-catalyzed click chemistry at a dye concentration of 1-5 μ M (final concentration) to guarantee for best signal-to-noise ratios, but substantially lower or higher concentrations can be used, if necessary. We successfully tested concentrations as low as 10 nM and as high as 50 μ M.

00:03:00

Note

The Hatzenpichler lab highly recommend dyes available at clickchemistrytools, in particular picolyl-based dyes. Theser provide a Cu-chelating motif which drastically improves reaction kinetics.

Cu(I) Click Chemistry (Immobilized)

15 If using immobilized biomass after dehydration, continue. Otherwise, skip to Step 18.

Add 12.5 μ L of each 100 mM sodium ascorbate and 100 mM aminoguanidine hydrochloride to 221 μ L PBS. Then, add the dye premix and invert the tube once (do not mix by vortex to maintain reducing conditions).

- Cover the sample with 20 μL of the click solution, transfer the slide into a humid chamber (water on tissue paper), and incubate in the dark at RT for 30 min. Increasing the incubation time is possible, but typically does not increase fluorescence signal.
 00:30:00
- 17 Wash the slide or filter three times for 3 min each in PBS-filled 50 mL tubes before dehydrating it by incubating it for 3 min in 50% EtOH at RT.

Cu(I) Click Chemistry (Solution)

18 If the biomass is in solution, all reagents (sodium ascorbate and aminoguanidine, followed after 3 min by the dye premix, final concentrations as described above) are added directly to the sample. Invert tubes once and incubate in the dark at RT for 30 min. Afterwards, wash samples three times with PBS and then one time in 50% EtOH (RT). Between washing steps, pellet samples via centrifugation for 5 min at 16,100g (or highest setting) at RT. Finally, resuspend biomass in a 1:1 mix of PBS:EtOH, transfer onto a glass slide, and air-dry

00:30:00

19