MAR 28, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.sfnebme

External link:
<https://doi.org/10.1371/journal.pone.0213493>

Protocol Citation: Jorge Carlos Ruiz Ruiz 2019. Determination of flavonoid content. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.sfnebme>

MANUSCRIPT CITATION:
 Dewanto V, Wu X, Adom KK, Liu RH. Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. J Agric Food Chem. 2002; 50: 3010-3014.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Aug 07, 2018

Determination of flavonoid content

PLOS One

Jorge Carlos Ruiz Ruiz¹

¹Universidad Anáhuac Mayab

Jorge Carlos Ruiz Ruiz

ABSTRACT

Flavonoid content determination using the aluminum chloride method.

MATERIALS

MATERIALS

Sodium nitrite **P212121**

water **Contributed by users**

aluminum chloride **Sigma Aldrich**

sodium carbonates **Sigma Aldrich**

catechin **Sigma Aldrich**

PROTOCOL integer ID:
14542

- 1** An aliquot of sample (0.1 mL) was mixed with 0.4 ml distilled water in a 1.5 mL microcentrifuge tube.
- 2** 0.03 mL of 5% NaNO₂ was added and the mixture was allowed to react for 5 min.
- 3** Following this, 0.03 mL of 10% AlCl₃ was added and the mixture stood for a further 5 min.
- 4** Finally, the reaction mixture was treated with 0.2 mL of 1 M Na₂CO₃ and 0.24 mL distilled water, and the absorbance at 510 nm.
- 5** Catechin (Sigma-Aldrich) was used as the standard and the results were expressed as µg/mL of catechin equivalents (CE).