OCT 25, 2019

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.8qwhvxe

External link:

https://animal.ifas.ufl.edu/hansen/lab_protocol_docs/detection_lipids_embryos_nile_red_fluorescence.pdf

Protocol Citation: Luciano Bonilla, Peter J Hansen 2019. Determining Lipid Content in Embryos using Nile Red Fluorescence. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.8qwhvxe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 25, 2019

Last Modified: Oct 25, 2019

PROTOCOL integer ID:
29174

Determining Lipid Content in Embryos using Nile Red Fluorescence

Luciano Bonilla¹, Peter J Hansen¹

¹University of Florida

University of Florida Bovine Embryo Lab

Tech. support email: hansen@animal.ufl.edu

Peter J Hansen

University of Florida

ABSTRACT

The following protocol is derived from the protocol described by Genicot et al. (Theriogenology 63: 1181- 1194, 2005) and is based on the fluorescence emitted by Nile Red when in association with lipid (see Fowler and Greenspan, J Histochem Cytochem 33: 833-836, 1985).

ATTACHMENTS

[detection_lipids_embryos_nile_red_fluorescence.pdf](#)

Keywords: preimplantation
embryo, Nile Red, lipid