

Oct 16, 2020

🌐 Development of an anatomic landmark-based measurement of the Achilles tendon

DOI

dx.doi.org/10.17504/protocols.io.beaejabe



Daniel Urness¹

¹Austin Thompson, Emelia Sodders, Erik Ensrud, James Meeker

Daniel Urness: [Oregon Health & Science University Department of Orthopaedics and Rehabilitation]



Daniel Urness

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.beaejabe>

Protocol Citation: Daniel Urness 2020. Development of an anatomic landmark-based measurement of the Achilles tendon.
protocols.io <https://dx.doi.org/10.17504/protocols.io.beaejabe>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it is working for us

Created: March 26, 2020

Last Modified: October 16, 2020

Protocol Integer ID: 34854

Keywords: Achilles tendon, Achilles tendon rupture, rupture, ultrasound, US, MRI, longer measurement of the achilles tendon, measurement of achilles tendon, achilles tendon measure, longer measurements of the achilles tendon, based achilles tendon measure, achilles tendon length, achilles tendon, achilles tendon introduction achilles tendon rupture, achilles tendon injury, tendon length, change in tendon length, degree of tendon, tendon, ultrasound measurement, accuracy of the anatomic landmark, ankle, slight external rotation with the ankle, manual palpation of anatomic landmark, flexion at the knee, using ultrasound, ultrasound technique, longer measurement, accurate measurement technique, measurement technique in the operating room, intraclass correlation coefficient between the anatomic landmark, measurement technique, development of an anatomic landmark, comparison with mri, anatomic landmark, lower leg, length from the medial head, conclusions while the anatomic landmark,

Abstract

Introduction

Achilles tendon ruptures (ATRs) occurred with an incidence of 2.5 per 100,000 person-years in 2016. This rate has been increasing over the last decade and it has been postulated that this is due to the increasing activity level of older people. With a high and increasing incidence of such a significant injury, further investigation must be done to optimize the treatment of ATRs. The degree of tendon lengthening has been correlated with clinical outcomes, with greater elongation being associated with worse outcomes. MRI and ultrasound techniques have been validated in measurement of Achilles tendon. We sought to develop a reliable, reproducible, and accurate measurement technique utilizing the manual palpation of anatomic landmarks that will be cost effective as well as convenient to perform, particularly intraoperatively.

Methods

Both lower legs of 10 healthy subjects without history of Achilles tendon injury were examined using ultrasound and anatomic landmark-based measurement techniques. Subjects sat upright on the exam table and legs were held in flexion at the knee with slight external rotation with the ankle held at 90°, allowing the lower leg to rest flat on the exam table. The length from the medial head of the gastrocnemius to the bottom of the non-compressed heel pad was measured by three raters using the ultrasound and anatomic landmark-based techniques for inter-rater reliability. The measurements were repeated one week later for intra-rater reliability. Ultrasound measurements had excellent inter-rater (0.93) and intra-rater (0.82) correlation coefficients, while good inter-rater (0.76) and intra-rater (0.86) correlation coefficients were observed among anatomic landmark-based measurements. Achilles tendon length measured with ultrasound and anatomic landmark-based techniques were compared using a paired t-test.

Results

The anatomic landmark-based technique produces longer measurements of the Achilles tendon (23.2 cm (sd 2.6 cm)) compared to measurements made using ultrasound (22.4 cm (sd 2.6)) ($p < 0.001$). On average, the anatomic landmark-based technique measures 0.8 cm (95% Confidence Interval: 0.4, 1.2) longer than the ultrasound technique. The intraclass correlation coefficient between the anatomic landmark-based and ultrasound measurements is 0.90.

Conclusions

While the anatomic landmark-based technique produces a longer measurement of the Achilles tendon, it may still be a reproducible measurement tool. If the change in tendon length is of interest, this technique may be a valid and simple way to monitor that variable. Comparison with MRI may be warranted to better determine the accuracy of the anatomic landmark-based Achilles tendon measure. These results compared with MRI may set the stage for further evaluation of this measurement technique in the operating room in subjects undergoing ATR repair.



Materials

Industrial steel framing square

Tape measure

Marking pen

Paper clip


alcohol pads

Ultrasound machine with gel

MRI machine

Troubleshooting

Safety warnings

 None

Positioning

- 1
 1. Position subject sitting upright with hip flexed, abducted, and externally rotated so the lateral aspect of the lower leg lays flat over the outline of the framing square
 2. Align the plantar aspect of the foot with the base of the framing square and the posterior aspect of the calf with the vertical portion of the square to make a 90 degree angle at the ankle joint.



Anatomic landmark-based measurement


- 2
 1. With subject positioned as above, palpate the step off where the medial head of the gastrocnemius becomes the Achilles tendon.



2. Mark above location with marking pen.
3. Using a tape measure, measure from that mark to the base of the non-compressed heel pad and record
4. Remove marking pen with alcohol pad and repeat for 3 measurements on both lower legs for a total of 6 measurements.
5. Rotate subjects to next rater until each rater has recorded measurements for each subject.

Ultrasound measurement

- 3
 1. With the subject positioned identically to the anatomic landmark-based technique, use the ultrasound to identify the confluence of the medial head of the gastrocnemius and the Achilles tendon.



2. Mark this landmark. (The authors used a paper clip to mark this location on the ultrasound and then marked the area with a marking pen)
3. Using a tape measure, measure from that mark to the base of the non-compressed heel pad and record.
4. Remove marking pen with alcohol pad and repeat for 3 measurements on both lower legs for a total of 6 measurements.
5. Rotate subjects to next rater until each rater has recorded measurements for each subject.

MRI Positioning

- 4
 1. Position patient supine with bilateral ankles at 90 degrees in MRI scanner.
 2. Image must capture medial head of the gastrocnemius and distalmost portion of the non-compressed heelpad.
 3. Obtain axial, coronal, and sagittal T1 series for bilateral legs.

MRI Measurement

- 5
 1. Identify confluence of the medial head of the gastrocnemius with the Achilles tendon using an axial cut and identify corresponding landmark on a sagittal or coronal image.
 2. Using either coronal or sagittal cut, identify distalmost portion of the non-compressed heelpad.
 3. Measure the distance between the two above landmarks using imaging software.
 4. Each measurement is made once on each leg by two separate raters.