MAY 17, 2020

DH5 α bacteria transformation protocol

Yoan Coudert¹¹Ecole Normale Supérieure de Lyon

CoudertLab

Yoan Coudert

Ecole Normale Supérieure de Lyon

OPEN ACCESS**DOI:**dx.doi.org/10.17504/protocols.io.bghijt4e**Protocol Citation:** Yoan Coudert 2020. DH5 α bacteria transformation protocol.**protocols.io**<https://dx.doi.org/10.17504/protocols.io.bghijt4e>**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited**Protocol status:** Working

We use this protocol and it's working

Created: May 17, 2020**Last Modified:** May 17, 2020**PROTOCOL integer ID:**

37130

1 Thaw bacteria on ice

- 2 Dispense 50µl in a microtube
- 3 Add 1 to 5 µl of DNA, and carefully flick the tube to mix cells and DNA
- 4 Place the mixture on ice for 10 minutes
- 5 Heat shock at 42°C for 30 seconds
- 6 Place on ice for 5 minutes
- 7 Pipette 450 µl of room temperature SOC (or LB) into the mixture
- 8 Incubate at 37°C for 1 hour. Shake vigorously at 250 rpm.
- 9 Warm selection plates at 37°C
- 10 Spread desired volume (150-350 µl) onto selection plate

- 11 Incubate overnight at 37°C
- 12 Store at 4°C after incubation of needed