JAN 03, 2018

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.kbacsie](https://doi.org/10.17504/protocols.io.kbacsie)

External link:
<https://doi.org/10.1371/journal.pone.0190595>

Protocol Citation: Bianca Hemmeryckx, Dries Bauters, H. Roger Lijnen 2018. Diet-induced obesity murine model. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.kbacsie>

MANUSCRIPT CITATION:
 Bauters D, Bedossa P, Lijnen HR, Hemmeryckx B (2018) Functional role of ADAMTS5 in adiposity and metabolic health. PLoS ONE 13(1): e0190595. doi: [10.1371/journal.pone.0190595](https://doi.org/10.1371/journal.pone.0190595)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Oct 15, 2017

Diet-induced obesity murine model

PLOS Biology

Bianca Hemmeryckx¹, Dries Bauters¹, H. Roger Lijnen¹

¹KU Leuven

Bianca Hemmeryckx

ABSTRACT

Obesity can be induced by exposing mice to a western-type diet high in sugar and fat for 15 weeks. Control animals receive a standard fat diet for 15 weeks.

GUIDELINES

-HFD is stored at +4°C

-Mouse needs to be out of cage, relaxed on a bench, a scalpel is used to cut of 1 mm of tail, the tail starts to bleed, glucose levels are measured on the second drop of blood.

MATERIALS

STEP MATERIALS

- ssniff® EF R/M acc. TD88137 mod **Contributed by users Catalog #E15721-3**
- Accu-Chek Aviva glucose strips **Contributed by users Catalog #06453970**
- Nembutal 60 mg/ml **Contributed by users**
- Dolethal 200 mg/ml **Contributed by users**
- Microflex Infusion set, orange **Contributed by users Catalog #240.05**
- Virkon™ S 50 tablets/bottle **Fisher Scientific Catalog #NC9821357**
- ssniff® EF R/M acc. TD88137 mod **Contributed by users Catalog #E15721-3**
- Accu-Chek Aviva glucose strips **Contributed by users Catalog #06453970**
- Nembutal 60 mg/ml **Contributed by users**
- Dolethal 200 mg/ml **Contributed by users**
- Microflex Infusion set, orange **Contributed by users Catalog #240.05**
- Virkon™ S 50 tablets/bottle **Fisher Scientific Catalog #NC9821357**

Last Modified: Feb 21, 2018

PROTOCOL integer ID:
8258

Keywords: obesity, adipose
tissue, murine model,
western-type diet

- 1 Randomize 8-weeks old male mice (C57BL/6J, C57BL/6N or gene-deficient mice with control littermates) to two groups based on body weight.
- 2 Give one group 7g standard chow (SFD, 10.9 kJ/g) each day/mouse, and the other group 7g western high-fat diet (HFD, 22 kJ/g; kcal from fat 42%, 43% from carbohydrates and 15% from protein) for 15 weeks. As food intake averages 4g/day, the animals are fed ad libitum. Provide drinking water ad libitum.

ssniff® EF R/M acc. TD88137 mod **Contributed by users Catalog #E15721-34**
- 3 House the animals in a temperature (20-23°C) and humidity (30-40%)-controlled room with a 12-hour light/dark cycle (7AM/7PM). Give 1-2 small wooden sticks to every cage containing mice on a HFD in order to allow correct wearing off of their teeth.
- 4 Replace drinking water and change the cages every week. Check the health of the animals every day.
- 5 Replace each diet weekly, record food intake and body weight weekly.
- 6 After 15 weeks of diet administration, fast animals for 6 hours (7:30 AM – 1:30 PM).
- 7 Determine blood glucose levels on a drop of blood taken from the nicked tail of a free-moving mouse using the Accu-Chek Aviva meter and Accu-Chek Aviva glucose strips.

Accu-Chek Aviva glucose strips **Contributed by users Catalog #06453970**

8 Sacrifice animals by intraperitoneal injection of 60 mg/kg sodium pentobarbital (Nembutal) or Dolethal.

-Nembutal (60 mg/ml): we make 10x dilution with saline.

-Dolethal (200 mg/ml): we make 30x dilution with saline.

Nembutal 60 mg/ml **Contributed by users**

Dolethal 200 mg/ml **Contributed by users**

9 When animals are asleep, collect blood from the retro-orbital sinus on trisodium citrate and processed for plasma (<http://dx.doi.org/10.17504/protocols.io.j9dcr26>). Store plasma in aliquots at -80°C.

10 Immobilize the animals, cut open the chest and wash out the heart and the entire circulatory system with saline by putting a perfusion needle in the apex (left ventricle). Make a small cut in the right atrium to allow blood to flow out.

Microflex Infusion set, orange **Contributed by users Catalog #240.05**

11 Make a midline incision in the skin and make a incision at the height of the hind limbs.

12 Pull skin to the side so the inguinal subcutaneous white adipose tissue (sWAT) depots are visible. Remove the internal lymph nodes. Dissect out the sWAT depots.

13 Make a midline incision in the muscle layer to open the abdominal cavity.

14 Locate the intra-abdominal gonadal (GON) WAT depots and liver. Removed them.

15 Flip the mouse, locate the interscapular brown adipose tissue depot and remove it.

- 16 Weigh all dissected organs.
- 17 Make portions of each tissue for RNA, protein extraction or histology.
- 18 Snap-freeze the portions for RNA and protein extraction and store at -80°C.
- 19 Process the portions for histology (see separate protocol).
- 20 Clean dissection material and bench with Vircon S, detergent, water and 70% ethanol.
Virkon™ S 50 tablets/bottle **Fisher Scientific Catalog #NC9821357**
- 21 Dispose of the animal body according to internal institutional guidelines.