OCT 14, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.76zhrf6

Protocol Citation: Igem Dusseldorf 2019. Disruption of Synechocystis cells.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.76zhrf6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 14, 2019

Last Modified: Oct 14, 2019

PROTOCOL integer ID:
 28601

🌐 Disruption of Synechocystis cells

Igem Dusseldorf¹

¹Heinrich-Heine Universität Düsseldorf

Igem Dusseldorf

ABSTRACT

Protocol for disrupting Synechocystis cells to get soluble and non-soluble protein in different fractions

- Carry out all steps on ice
- Thylacoid-buffer (TP)
- 50 mM HEPES/NaOH
- 5 mM MgCl₂
- 50 mM EDTA. pH 8
- 1 Tablet Protease inhibitor (Roche complete ULTRA tablets) per 10 ml TP

Cell disruption

Add 200 µl TP to a pellet of 20 ml Cell culture of OD₇₅₀ ~0,5 and resuspend the pellet

Add ca. 200 µl glas beads (1:1 mix of 0,1mm beads and 0,25 mm beads)

Use program 2 of the precellys to disrupt the cells at room temperature, put on ice immediately afterwards

Centrifuge for 5 min at 4°C and 4000 rpm to seperate glas beads and cellular waste from protein

Take supernatant (Whole cell extract 1)

Wash glas beads with 200 µl TP and repeat Precellys disruption

Centrifuge for 5 min at 4°C and 4000 rpm to separate glas beads and cellular waste from protein

Take supernatant (Whole cell extract 2)

Mix Whole cell extract 1 and 2

Optional: Centrifuge for 45 min at 21000 rpm and 4°C to seperate soluble from non-soluble protein. Soluble protein will contain Phycocyanin, non-soluble protien will contain Chlorophyll.

Freeze the protein until use, keep on ice when using it.