

Dec 24, 2019 Version 3

Dissection and immunohistochemistry of mouse vagal ganglia V.3

DOI

dx.doi.org/10.17504/protocols.io.baumieu6

Thomas Taylor-Clark¹, Seol-Hee Kim¹

¹University of South Florida

SPARC

Tech. support email: info@neuinfo.org



Seol-Hee Kim

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.baumieu6

Protocol Citation: Thomas Taylor-Clark, Seol-Hee Kim 2019. Dissection and immunohistochemistry of mouse vagal ganglia. [protocols.io](https://dx.doi.org/10.17504/protocols.io.baumieu6) <https://dx.doi.org/10.17504/protocols.io.baumieu6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: December 24, 2019

Last Modified: December 24, 2019

Protocol Integer ID: 31341

Keywords: vagal ganglia, immunohistochemistry, nodose, jugular



Abstract

Mice are euthanized, perfused with fixative and the vagal ganglia extracted. Vagal ganglia are then cyosectioned. Slices are stained for protein expression using immunohistochemistry. Expression of specific proteins and reporter proteins isarevisualized using microscopy.

Materials

MATERIALS

⊗ Triton X-100 **Bio-Rad Laboratories Catalog #1610407**

⊗ Bovine Serum Albumin (BSA) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #A7906**

⊗ DPX **Merck Millipore (EMD Millipore) Catalog #1.00579.0500**

⊗ Sucrose

⊗ Tween 20 **Merck MilliporeSigma (Sigma-Aldrich) Catalog #P1379**

⊗ Paraformaldehyde **Merck MilliporeSigma (Sigma-Aldrich) Catalog #P6148**

⊗ Phosphate buffered saline powder, pH 7.4, for preparing 1 L solutions **Merck MilliporeSigma (Sigma-Aldrich) Catalog #P3813**

⊗ alexa fluor 647 chicken anti-goat **Invitrogen - Thermo Fisher Catalog #A21469**

- 1 6 to 8 weeks old mice (male) were euthanized by CO₂ inhalation.
- 2 Mice were transcardially perfused with phosphate buffered saline (PBS) to remove blood followed by 3.7% formaldehyde (PFA) in PBS.
- 3 Vagal ganglia (including both jugular and nodose ganglia) were collected and fixed with 3.7% PFA for 2 hours on ice.
- 4 Vagal ganglia were transferred to 18% sucrose solution for overnight at 4 °C.
- 5 Tissue were mounted in optimal cutting temperature compound and frozen
- 6 Tissue were cut at 20 µm thickness and collected onto superfrost plus microscopy slide glass
- 7 Tissue were washed with PBS three times for 10 min.
- 8 Tissue were blocked with blocking buffer (1% bovine serum albumin/10% normal donkey serum/0.3% triton x-100 in PBS) for 45 min at room temperature.
- 9 Tissue were incubated primary antibodies diluted in blocking buffer overnight at 4 °C.

Set 1. Rabbit anti-TrkA (1:300, 06-574, Millipore) and Goat anti-VR1 (1:150, SC-12498, Santa Cruz)
Set 2. Rabbit anti-TrkA (1:300, 06-574, Millipore) and Goat anti-TrkB (1:300, AF1494, R&D Systems)
- 10 Tissue were washed with 0.2% tween20 in PBS (PBST) for 10 min for three times at room temperature.
- 11 Secondary antibodies were diluted in the 1% bovine serum albumin/5% normal donkey serum/0.2% PBST and tissue were incubated with it for 1 hour at room temperature.



1. Alexa Fluor 488 donkey anti-rabbit (1:300, A21206, Invitrogen)
2. Alexa Fluor 647 Chicken anti-Goat (1:300, A21469, Invitrogen)

- 12 Tissue were washed with 0.2% tween20 in PBS (PBST) for 10 min for three times at room temperature.
- 13 Tissue were cover-slipped with DPX for imaging and left to dry in the dark overnight.