OCT 14, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.766hrhe

Protocol Citation: Igem Dusseldorf 2019. DNA Clean & Concentrator . **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.766hrhe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 14, 2019

Last Modified: Oct 14, 2019

PROTOCOL integer ID:
 28606

🌐 DNA Clean & Concentrator

Igem Dusseldorf¹

¹Heinrich-Heine Universität Düsseldorf

Igem Dusseldorf

ABSTRACT

- Plasmid DNA
- Binding Buffer
- DNA Wash Buffer
- DNA Elution Buffer or

Preparation

Add 24 ml 100% ethanol (26 ml 95% ethanol) to the 6 ml DNA Wash Buffer concentrate. Add 96 ml 100% ethanol (104 ml 95% ethanol) to the 24 ml DNA Wash Buffer concentrate.

Perform all centrifugation at $\geq 10000 \times g$.

Implementation

In a 1,5 ml microcentrifuge tube, add 2 - 7 volumes of DNA Bindung Buffer to each volume of DNA sample. Mix briefly by vortexing.

Application	Bindung Buffer : sample
plasmid, genomic DNA (> 2 kb)	2 : 1
PCR product, DNA fragment	5 : 1
ssDNA	7 : 1

Transfer mixture to a Zymo- Spin Column in a Collection Tube.

Centrifuge for 30 seconds. Discard the flow-through.

Add 200 µl DNA Wash Buffer to the column. Centrifuge for 30 seconds. Repeat the wash step.

Add $\geq 6\mu\text{l}$ DNA Elution Buffer or water directly to the column matrix and incubate at room temperature for one minute. Transfer the column to a 1,5 ml microcentrifuge tube and centrifuge for 30 seconds to elute the DNA.