MAY 22, 2018

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.j9gcr3w](https://doi.org/10.17504/protocols.io.j9gcr3w)

External link:

<https://doi.org/10.1371/journal.pone.0196204>

Protocol Citation: Michiyo Tsuru 2018. DNA exon and SNP analyses. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.j9gcr3w>

MANUSCRIPT CITATION:

Tsuru M, Ono A, Umeyama H, Takeuchi M, Nagata K (2018) Ubiquitin-dependent proteolysis of CXCL7 leads to posterior longitudinal ligament ossification. PLoS ONE 13(5): e0196204. doi: [10.1371/journal.pone.0196204](https://doi.org/10.1371/journal.pone.0196204)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Oct 11, 2017

DNA exon and SNP analyses

PLOS One

Michiyo Tsuru¹

¹Clinical Proteomics and Gene Therapy Laboratory, Kurume University, Fukuoka, Japan

Clinical Proteomics and Gene Therapy Laboratory

Michiyo Tsuru

Kurume University, Clinical Proteomics and Gene Therapy

ABSTRACT

Genomic DNA was extracted from white blood cells (PAXgene-RNA tubes) using the PAXgene blood RNA kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions (PAXgene blood RNA kit handbook version 2, April 2008). Target gene and adjacent DNA exons and SNP analyses were conducted using PRISM3730 DNA Analyzer (Applied Biosystems, Foster City, CA, USA) and TaqMan SNP Genotyping Assay (Applied Biosystems) according to the manufacturer's instructions. (Hokkaido System Science, Sapporo, Japan).

ATTACHMENTS

[exon2.pdf](#) [exon3.pdf](#) [exon1.pdf](#)

Last Modified: May 22,
2018

PROTOCOL integer ID:
8200