VERSION 1
NOV 23, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bcmriu56

Protocol Citation: Lilan Hao
2020. DNA extraction for
human microbe samples..
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bcmriu56>

License: This is an open
access protocol distributed
under the terms of
the [Creative Commons
Attribution License](#), which
permits unrestricted use,
distribution, and reproduction
in any medium, provided the
original author and source are
credited

Protocol status: Working
We use this protocol and it's
working

Created: Feb 17, 2020

Last Modified: Nov 25, 2020

PROTOCOL integer ID:
33169

DNA extraction for human microbe samples. V.1

In 8 collections

Lilan Hao¹

¹BGI-Shenzhen, Shenzhen 518083, China.

GigaScience Press

BGI

Lilan Hao

ABSTRACT

This protocol is used to clarify the process of total DNA extration for human microbe samples.

1 200µl samples were mixed with 500 µl TE buffer and 20 µl of 10 mg/ml lysozyme solution.

- 2 Incubate at 37 °C for 90min. Invert the tube several times to mix every 15 minutes. Centrifuge at 13,000 rcf/min for 10 min at 4 °C and discard the supernatant.
- 3 Add 300 µl of 20 mg/ml proteinase K and 5µl of 10% SDS. Vortex to resuspend cells.
- 4 Incubate at 55 °C in a shaking water bath for 120 min to effect complete lysis. Invert the tube several times to mix every 15 minutes.
- 5 Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then transfer the supernatant to a nuclease-free 2 ml microfuge tube.
- 6 Add 700 µl phenol-chloroform-isoamylalcohol (25:24:1) and vortex to mix well, centrifuge at 13,000 rcf/min for 10 min at 4 °C, then transfer the supernatant to a nuclease-free 2 ml microfuge tube.
- 7 Add 700 µl chloroform-isoamylalcohol (24:1) and vortex to mix well. Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then transfer the supernatant to a nuclease-free 1.5 ml microfuge tube.
- 8 Add 1/10 volume of 3M sodium acetate, 2 µl of 5 mg/ml glycogen and 800 µl cold isopropanol. Incubate at -20 °C for the night.
- 9 Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then discard the supernatant.

- 10** Washed with 800 µl of 70% ethanol. Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then discard the supernatant.
- 11** Wash the precipitate with a second 800 µl of 70% ethanol and centrifuge as above. discard the supernatant.
- 12** Let sit for 15 min to dry the precipitate. Add 1×TE buffer to resuspend the precipitate.
- 13** Add 1µl RNase A to samples and invert tube several times to mix. Incubate at 37 °C for 30 minutes.