

Nov 23, 2020 Version 1

DNA extraction for human microbe samples. V.1

 [GigaByte](#)

 Peer-reviewed method


 In 8 collections

DOI

dx.doi.org/10.17504/protocols.io.bcmriu56

Lilan Hao¹

¹BGI-Shenzhen, Shenzhen 518083, China.

 GigaScience Press

 BGI



Lilan Hao

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bcmriu56

External link: <https://doi.org/10.46471/gigabyte.9>

Protocol Citation: Lilan Hao 2020. DNA extraction for human microbe samples.. [protocols.io](#)

<https://dx.doi.org/10.17504/protocols.io.bcmriu56>

Manuscript citation:

Chen C, Hao L, Wei W, Li F, Song L, Zhang X, Dai J, Jie Z, Li J, Song X, Wang Z, Zhang Z, Zeng L, Du H, Tang H, Zhang T, Yang H, Wang J, Brix S, Kristiansen K, Xu X, Wu R, Jia H, The female urinary microbiota in relation to the reproductive tract microbiota. GigaByte. 2020.

<https://doi.org/10.46471/gigabyte.9>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: February 17, 2020

Last Modified: November 24, 2020

Protocol Integer ID: 33169

Abstract

This protocol is used to clarify the process of total DNA extraction for human microbe samples.

- 1 200 μ l samples were mixed with 500 μ l TE buffer and 20 μ l of 10 mg/ml lysozyme solution.
- 2 Incubate at 37 °C for 90min. Invert the tube several times to mix every 15 minutes. Centrifuge at 13,000 rcf/min for 10 min at 4 °C and discard the supernatant.
- 3 Add 300 μ l of 20 mg/ml proteinase K and 5 μ l of 10% SDS. Vortex to resuspend cells.
- 4 Incubate at 55 °C in a shaking water bath for 120 min to effect complete lysis. Invert the tube several times to mix every 15 minutes.
- 5 Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then transfer the supernatant to a nuclease-free 2 ml microfuge tube.
- 6 Add 700 μ l phenol-chloroform-isoamylalcohol (25:24:1) and vortex to mix well, centrifuge at 13,000 rcf/min for 10 min at 4 °C, then transfer the supernatant to a nuclease-free 2 ml microfuge tube.
- 7 Add 700 μ l chloroform-isoamylalcohol (24:1) and vortex to mix well. Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then transfer the supernatant to a nuclease-free 1.5 ml microfuge tube.
- 8 Add 1/10 volume of 3M sodium acetate, 2 μ l of 5 mg/ml glycogen and 800 μ l cold isopropanol. Incubate at -20 °C for the night.
- 9 Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then discard the supernatant.
- 10 Washed with 800 μ l of 70% ethanol. Centrifuge at 13,000 rcf/min for 10 min at 4 °C, then discard the supernatant.
- 11 Wash the precipitate with a second 800 μ l of 70% ethanol and centrifuge as above. discard the supernatant.
- 12 Let sit for 15 min to dry the precipitate. Add 1 \times TE buffer to resuspend the precipitate.

-
- 13 Add 1 μ l RNase A to samples and invert tube several times to mix. Incubate at 37 °C for 30 minutes.