OCT 25, 2020

DNT Detection In Soil

Zhujun Wei¹

¹2020 iGEM NEFU China

2020 iGEM NEFU China

Zhujun Wei

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bnxtmfnn

Protocol Citation: Zhujun Wei 2020. DNT Detection In Soil. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bnxtmfnn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 25, 2020

Last Modified: Oct 25, 2020

PROTOCOL integer ID:
43731

- 1 The 1% bacteria (PYB1s-yqjF4-luxCDABE) were cultured in LB containing 0.1% streptomycin (50 mg/mL) for 24 h with aeration at 37°C to reach a final density of 1×10^8 cells/mL.

- 2 Preparation of 2% Alginate solution, 1% calcium-chloride solution, experimental soil: dissolve 2g Alginate in 100 mL distilled water (2% w/w) and 1g calcium chloride in 100 mL distilled water (1% w/w). The experimental soil was filtrated prior to use.
- 3 Sterilize the 2% Alginate solution, 1% calcium-chloride solution, (121°C, 15min), and experimental soil (121°C, 1.5h).
- 4 Add 0, 0.25, 0.5 and 1mg of DNT powder (0, 250, 500 and 1000mg/kg) to the wells of a 12-well plate. Take 1g of experimental soil and drop it onto the DNT in the 12-well plate on a clean bench.
- 5 The engineered bacteria were added into 2% Alginate solution at a 1:9 (v/v) ratio.
- 6 Drop 200 µL of 1% (w/w) calcium-chloride solution into the 1mL alginate-bacterial suspension droplets (the volume ratio between the Alginate mixture and calcium chloride solution was 5:1). Leave the mix for 1 minute.
- 7 The mixed Alginate gel was tiled on the soil.