

Oct 25, 2020

DNT Induction

DOI

dx.doi.org/10.17504/protocols.io.bnxsmfne

Zhujun Wei¹

¹2020 iGEM NEFU China

2020 iGEM NEFU China



Zhujun Wei

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bnxsmfne

Protocol Citation: Zhujun Wei 2020. DNT Induction. protocols.io <https://dx.doi.org/10.17504/protocols.io.bnxsmfne>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 25, 2020

Last Modified: October 25, 2020

Protocol Integer ID: 43730



- 1 The DNT powder (>99%) was dissolved in 80% acetonitrile solution, prepared as 10g/L stock solution, and stored at 4°C.
- 2 The glycerin strains were first grown overnight in LB medium at 37°C supplied with appropriate antibiotics.
- 3 The bacteria cultured overnight were diluted by 1/100 times into fresh LB medium, and a certain amount of DNT stock was added to make DNT reach an expected concentration. Then, the samples (200 µL) were added to the wells of a clear 96-well plate with LB medium only as a negative control.
- 4 The RFU (relative fluorescence units) and OD₆₀₀ were measured every 30 minutes using the multi-functional enzyme reader at 37°C shaking culture condition, for a period of 12h. (For RFU measurement, 200 µL of LB was used for calibration, with an excitation wavelength 485nm, an excitation bandwidth 20nm, an emission wavelength 535nm, and an emission width 70nm)
- 5 The experiment was repeated three times for each sample.