


Sep 24, 2019

## Domestication of L0 parts for Loop type IIS (Bsal and Sapl)

DOI

[dx.doi.org/10.17504/protocols.io.4whgxb6](https://dx.doi.org/10.17504/protocols.io.4whgxb6)



Eftychis Frangedakis<sup>1</sup>, Marta Marta Tomaselli<sup>2</sup>, Marius Rebmann<sup>3</sup>, Susana Sauret-Gueto<sup>3</sup>

<sup>1</sup>University of Cambridge; <sup>2</sup>University of Cambridge, Open Plant;

<sup>3</sup>Plant Sciences, University of Cambridge, OpenPlant

OpenPlant Project



Eftychis Frangedakis

University of Cambridge, Plant Sciences

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.4whgxb6](https://dx.doi.org/10.17504/protocols.io.4whgxb6)

**Protocol Citation:** Eftychis Frangedakis, Marta Marta Tomaselli, Marius Rebmann, Susana Sauret-Gueto 2019. Domestication of L0 parts for Loop type IIS (Bsal and Sapl). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.4whgxb6>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

We use this protocol and it's working

**Created:** June 28, 2019

**Last Modified:** September 24, 2019

**Protocol Integer ID:** 25257

- 1 Domestication of internal Bsal and Sapl sites is necessary when a DNA sequence of interest contains an internal Bsal or Sapl recognition site. For domestication, overlapping PCR can be used. Briefly, the sequence can be amplified as two separate PCR fragments upstream (fragment 1) and downstream (fragment 2) the recognition site. The enzyme recognition site will form part of the reverse primer of fragment 1 and the forward primer of fragment 2, that will be specially designed with a single nucleotide mismatch to alter the recognition site (taking care to not alter amino acid composition if the region is protein coding). Amplification primers will also contain Sapl recognition sites that will allow the two fragments to be ligated together into the pUAP-pe. For example if you need to domesticate the following sequence:

5'-atggcgtcaattagtggatgcgcagctgtgccttgagc..... *ctattccagatt**GGTCT-C**-agccaatgcaactgaagggc* .....*gattatcacagagggtatggaaggactataactaa*-3'

(fragment 1: underlined sequence, Bsal recognition sequence: sequence in bold capital letters and fragment 2: italicized sequence), the primers to be designed are

Primer1: 5'-ga**GCTCTTC**gtctcgaatggcgtcaattagtggatgc-3',

Primer2: 5'-ta**GCTCTTC**gCacc-aatctggaatag -3',

Primer3: 5'-ga**GCTCTTC**gtc**Gc**-agccaatg.caactgaagggc-3', and

Primer4: 5'-ta**GCTCTTC**gtctcaaagcttagtatatagtccatc-3'

(Sapl recognition sequence: sequence in bold capital letters, common syntax overhangs: underlined sequence and primer mismatch: sequence in bold underlined capital letter). Primer1 and Primer4 are designed according to Level 0 parts primer design.

2

