OCT 14, 2019

🌐 Double Digestion of Insert DNA

Igem Dusseldorf¹

¹Heinrich-Heine Universität Düsseldorf

Igem Dusseldorf

ABSTRACT

Combined enzyme volume should not exceed 1/10 of the total reaction volume.

For **double** digestion with **two** FastDigest restriction enzymes, mix:

Write down the name DNA tube used and its concentration.

clean and concentrate the sample after digest in 10µL Water

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.76qhrdw

Protocol Citation: Igem Dusseldorf 2019. Double Digestion of Insert DNA. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.76qhrdw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 14, 2019

Last Modified: Oct 14, 2019

PROTOCOL integer ID:
28592

PCR Purified 2016.07.25	ng/uL
Gene DNA Tube X	50
Component	Volume x2µl]
Sterile MilliQ Water	24
10x FastDigest Buffer	4
PCR-amplified DNA	8
FastDigest Enzyme 1	2
FastDigest Enzyme 2	2
Total Volume	40

Total ng of DNA	400
-----------------------	-----

Mix gently and spin down.

Incubate at enzyme suitable conditions (45min)

Inactivate the enzyme at suitable conditions