FEB 19, 2019

🌐 Double picking protocol for tracking

Priota Islam¹

¹Imperial College London

Behavioural Genomics

Priota Islam

Imperial College London

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.yb2fsqe

Protocol Citation: Priota Islam 2019. Double picking protocol for tracking.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.yb2fsqe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 19, 2019

Last Modified: Feb 19, 2019

PROTOCOL integer ID:
 20570

Imaging plates

- 1
 1. Prepare and pour low peptone NGM onto 35mm imaging plates
 2. Put them in the cold room for at least 2 days before use
 3. Seed imaging plates with 50ul of freshly made 1:10 solution (OP50:M9 solution) the day

before recording

4. Leave on bench to dry with lid on overnight

Day before recording

- 2
 1. Pick around 45 L4s with a hair pick onto seeded NGM plate from maintenance plates having a mixed population of worms
 2. Leave the plates in a 20C incubator for ~24hrs

Day of recording

- 3
 1. Using a hair pick – pick 5 young adults onto the seeded imaging plates
 2. Put the plates on the tracker – agar side up with the lids off
 3. Wait 30min for the worms to acclimatise
 4. Record for 15min (See protocol ***Imaging on the multiworm tracker*** for detailed tracking instructions)