VERSION 2

DEC 20, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.baq5idy6

Protocol Citation: Diep R Ganguly 2019. Dye-terminator DNA sequencing. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.baq5idy6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Dec 20, 2019

Last Modified: Dec 20, 2019

PROTOCOL integer ID:
 31229

Keywords: Sanger sequencing, Dye-terminator sequencing

Dye-terminator DNA sequencing V.2

Diep R Ganguly¹

¹University of Pennsylvania

Pogson Group

Diep R Ganguly

University of Pennsylvania, The Australian National Universi...

ABSTRACT

This protocol (based on the BigDye® Terminator v3.1 Cycle Sequencing Kit) is for performing terminator cycling sequencing reactions for Sanger sequencing of amplified PCR products or plasmid DNA on the 3130X genetic analyser (Applied Biosystems).

ATTACHMENTS

[BigDye Terminator v3.1.pdf](#)

MATERIALS

MATERIALS

- ⊗ Antarctic Phosphatase - 1,000 units **New England Biolabs Catalog #M0289S**
- ⊗ 96 well PCR Plate Non-skirted **Phenix Research Catalog #MPS-499**
- ⊗ Nuclease-free water (e.g. MilliQ or HPLC grade water) **Contributed by users**
- ⊗ primers **Contributed by users**
- ⊗ EDTA **Contributed by users**
- ⊗ 10 mM dNTPs **Life Technologies Catalog #10297-018**
- ⊗ Ethanol **Merck Millipore Catalog #100983**
- ⊗ BigDye™ Terminator v3.1 Cycle Sequencing Kit **Thermo Fisher Catalog #4337454**
- ⊗ Exonuclease I (E. coli) **NEB Catalog #M0293S**
- ⊗ Hi-Di™ Formamide **Thermo Fisher Scientific Catalog #4311320**

BEFORE START INSTRUCTIONS

Optimize PCR cycling (if sequencing amplified PCR products) to ensure your reaction produces a single product. Perform gel excision or PCR clean-up with the potential inclusion of incubating with Antarctic phosphatase and Exonuclease 1 to dephosphorylate and degrade unincorporated dNTPs in PCR reactions to prepare templates for DNA sequencing.

Terminator cycling reaction

- 1 Perform sequencing reaction in PCR tubes (or 96-well plate) with BigDye Terminator cycling kit and forward or reverse primers.

30m

Component	Volume (μl)
2.5X Reaction Ready Mix	1
5X BigDye Sequencing buffer	2
20 μM F/R Primer	0.5
Template (plasmid or cleaned PCR product)	150 ng dsDNA
Nuclease-free water	to 10 μl

BigDye Terminator Cycling reaction

5x reaction buffer: 400 mM TRIS, 10 mM MgCl₂

- 2 Run the following thermal cycling protocol:
- 1 min at 96 °C
 - 30-40 cycles: 96 °C for 10 seconds, 50 °C for 5 seconds, and 60 °C for 4 min.
 - Hold at 4-12 °C.

4h

Purification

1h 30m

- 3 Transfer PCR reaction to eppendorf tube. To the reaction, add 2.5 μL of 125 mM EDTA (make sure it touches bottom of tube).

- 4 Add 30 μ L of 100% ethanol, mix well (inversion).
- 5 Incubate at room temperature for 15 minutes.
- 6 Centrifuge at 4 °C at max speed for 30 minutes.
- 7 Discard supernatant and add 50 μ L of ice-cold 70% ethanol.
- 8 Centrifuge at 4 °C at max speed for 5 minutes.
- 9 Discard supernatant and allow to air-dry in the dark for >15 minutes.

Prepare for sequencing

- 10 Resuspend the pellet (likely transparent) in 7.5 μ L HiDi Formamide (add to any empty wells). Incubate at RT for 5 minutes then transfer to plate. Spin down briefly.
- 11 Incubate plate at 95 °C for 3 minutes (denature) then place immediately on ice. Spin down briefly.

-
- 12 Submit for sequencing on 3130X genetic analyser (Applied Biosystems). Keep samples on ice.