

Oct 05, 2025

E.coli DH10B Competent Cell Preparation

DOI

<https://dx.doi.org/10.17504/protocols.io.kxygx415dl8j/v1>

igemnthuth¹

¹iGEM NTHU_Taiwan

iGEM NTHU



igemnthuth

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.kxygx415dl8j/v1>

Protocol Citation: igemnthuth 2025. E.coli DH10B Competent Cell Preparation. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.kxygx415dl8j/v1>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working



Created: October 03, 2025

Last Modified: October 05, 2025

Protocol Integer ID: 228956

Keywords: competent cells suitable for electroporation, subsequent plasmid introduction, hosts for subsequent plasmid introduction, electroporation, plasmid, competent cell, cell, preparation

Abstract

Preparation of *E. coli* competent cells suitable for electroporation, to serve as hosts for subsequent plasmid introduction.

Materials

- Cultured log-phase DH10B cells (200 mL culture, OD600 $\approx$ 0.6–0.8)
- Ice-cold sterile 10% glycerol solution (filter-sterilized, pre-chilled at 4 °C or on ice, prepare at least 100 mL)
- Sterile centrifuge bottles (250–500 mL)
- Sterile microcentrifuge tubes (1.5 mL, 10 tubes)
- Sterile pipette tips and micropipettes
- Ice bucket and crushed ice bath (all procedures must be kept cold)
- Refrigerated centrifuge (set to 4 °C, 4500 $\times$ g)
- Shaking incubator (37 °C, 220 rpm)



E.coli DH10B Competent Cell Preparation

1h 10m

- 1 Obtain log-phase culture ($OD_{600} \approx 0.6-0.8$) and immediately place it on ice to cool for 10 minutes (gently swirling during cooling is recommended). 10m
- 2 Transfer 200 mL of culture into pre-chilled centrifuge bottles and centrifuge at $4500 \times g$ for 10 minutes at $4^\circ C$. 10m
- 3 Carefully discard the supernatant without disturbing the cell pellet. 5m
- 4 Add 50 mL of ice-cold 10% glycerol. Mix by repeatedly pipetting up and down until the suspension is evenly turbid. Centrifuge again at $4500 \times g$ for 10 minutes at $4^\circ C$. 10m
- 5 Repeat steps 3–4 for a total of two washes. 30m
- 6 After the final centrifugation, discard the supernatant and resuspend the entire pellet in 1 mL of ice-cold 10% glycerol. Transfer the suspension into sterile 1.5 mL tubes. 5m
- 7 Aliquot 50–100 μL per tube. Keep on ice for immediate use, or snap-freeze in liquid nitrogen and store at $-80^\circ C$ (stable for 1–2 weeks).