VERSION 3

MAR 01, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.yrpfv5n

Protocol Citation: Kenneth Schackart, Kattika Kaarj 2019. *E. coli* K12 DNA Extraction. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.yrpfv5n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Mar 01, 2019

Last Modified: Mar 01, 2019

PROTOCOL integer ID:
 21007

E. coli K12 DNA Extraction V.3

Kenneth Schackart¹, Kattika Kaarj¹

¹University of Arizona

Yoon Lab

Kenneth Schackart
 University of Arizona

ABSTRACT

How to extract DNA from *E. coli* K12 using Wizard® Genomic DNA Purification Kit by Promega®.

I do not claim any credit for the development of this protocol. It has been adapted from the protocol detailed in: Wizard Genomic DNA Purification.pdf .

MATERIALS

MATERIALS

Wizard(R) Genomic DNA Purification Kit **Promega Catalog #A1620**

Additional materials:

- 1.5 mL microcentrifuge tubes
- Isopropanol, room temperature
- 70% ethanol, room temperature
-

Culture bacteria

- 1 Culture *E. coli* K12 in BHI broth overnight.
2 mg lyophilized *E. coli* K12 in 10 mL BHI broth.

Pellet the cells

- 2 Add 1 mL cell suspension to 1.5 mL microcentrifuge tube.
- 3 Centrifuge at $13,000\text{--}16,000 \times g$ for 00:02:00 .
- 4 Remove supernatant.

Lyse nuclei

- 5 Add 600 μL of Nuclei Lysis Solution.
- 6 Gently pipet until the cells are resuspended.
- 7 Incubate at 80 °C on heating block for 00:05:00 to lyse the cells.
- 8 Cool to room temperature.

Degrade RNA

- 9 Add 3 μL RNase Solution to the cell lysate.
- 10 Invert 2-5 times to mix.
- 11 Incubate at 37 °C for 00:15:00 to 01:00:00 .
- 12 Cool to room temperature.

Precipitate proteins

- 13 Add 200 μL of Protein Precipitation Solution to the RNase-treated cell lysate.
- 14 Vortex vigorously at high speed for 00:00:20 .
- 15 Incubate on ice for 00:05:00

16 Centrifuge at 13,000-16,000 × *g* for 00:03:00 .

Harvest DNA

17 Transfer the supernatant containing the DNA to a clean 1.5 mL microcentrifuge tube containing 600 µL isopropanol.

Note

Some supernatant may remain in the original tube containing the protein pellet. Leave this residual to avoid contaminating the DNA solution with the precipitated protein.

18 Gently mix by inversion until the thread-like strands of DNA form a visible mass.

Wash and dry DNA

19 Centrifuge at 13,000-16,000 × *g* for 00:02:00 .

20 Carefully pour off the supernatant and drain the tube on clean absorbent paper.

21 Add 600 µL of 70% ethanol and gently invert the tube several times to wash the DNA pellet.

22 Centrifuge at 13,000-16,000 × *g* for 00:02:00 .

23 Carefully aspirate the ethanol.

24 Drain the tube on clean absorbent paper and allow to air-dry for 10-15 minutes.

Rehydrate DNA

25 Add 100 µL of DNA rehydration solution to the tube.

26 Rehydrate by incubating the solution overnight at room temperature or 4 °C .

27 Store DNA at 2 °C to 8 °C .