

Mar 28, 2019

Version 1

© E gene amplification V.1

DO

dx.doi.org/10.17504/protocols.io.zkjf4un

Bo Yi¹, Yi Chen¹, Xiao Ma¹, Haibin Wang², Rong Wang¹, Keqin Ding¹, Lei Xie¹, Dongliang Zhang¹, Shuli Jiao¹, Xuying Lao¹, Yi-Chen Chiang³, Yanhua Su³, Benhua Zhao³, Guozhang Xu¹, Tianmu Chen³

¹Ningbo Municipal Center for Disease Control and Prevention;

³State Key Laboratory of Molecular Vaccinology and Molecular Diagnostics, School of Public Health, Xiamen University



Tianmu Chen

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account





DOI: https://dx.doi.org/10.17504/protocols.io.zkjf4un

Protocol Citation: Bo Yi, Yi Chen, Xiao Ma, Haibin Wang, Rong Wang, Keqin Ding, Lei Xie, Dongliang Zhang, Shuli Jiao, Xuying Lao, Yi-Chen Chiang, Yanhua Su, Benhua Zhao, Guozhang Xu, Tianmu Chen 2019. E gene amplification. **protocols.io** https://dx.doi.org/10.17504/protocols.io.zkjf4un

²Haishu District Center for Disease Control and Prevention;



License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: March 28, 2019

Last Modified: March 28, 2019

Protocol Integer ID: 21867

Keywords: amplification, gene

Troubleshooting



- 1 The dengue virus strains obtained from cell culture were extracted by TGuide S32 magnetic bead method DNA/RNA extraction kit of Tiangen Biochemical Technology (Beijing) Co., Ltd. and amplified by one step RNA PCR Kit (Code No: DRR057A) reagent of Bao Bioengineering (Dalian) Co., Ltd.
- 2 The full sequence of E gene amplified by primer sequence was D1 (E) F: CAA GAA CCG AAA CA / GT GGA TGT C; D1 (E) R: GGC TGA TCG AAT TCC ACA CAC, and the length of amplified fragment was 1849 bp.
- 3 Reaction conditions: reverse transcription at 50 °C 30 min, reverse transcription at 94°C 2 min, reverse transcription at 94°C 30 s, reverse transcription at 52°C 30 s, reverse transcription at 72°C 2 min, 40 cycles and extension at 72°C. The amplified product was 5 micron L, and 1.5% agarose gel electrophoresis was used to confirm the reaction product according to Marker position.